



Digitized by the Internet Archive
in 2019 with funding from
University of Alberta Libraries

<https://archive.org/details/Smithers1965>



Thesis
1965
77

TELLURIUM ISOTOPE FRACTIONATION
IN NATURE AND IN THE
LABORATORY

by

Ronald M. Smithers

A THESIS
SUBMITTED TO THE FACULTY OF GRADUATE STUDIES
IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE
OF MASTER OF SCIENCE

DEPARTMENT OF PHYSICS
EDMONTON, ALBERTA
APRIL, 1965

ABSTRACT

It is well known that natural and laboratory processes can alter isotopic abundances. This thesis is an initial study to find the extent to which tellurium isotope ratios may be altered.

In the introduction, a brief review of historical isotope studies is given and some of the natural mechanisms which alter isotope abundances summarized. Isotope fractionation studies of sulphur and selenium are reviewed more extensively because tellurium is similar to sulphur and selenium.

The vibrational frequencies are computed for isotopic compounds of tellurium. From these, vibrational partition function ratios are calculated and these ratios are in turn used to calculate equilibrium constants for various $\text{Te}^{130} - \text{Te}^{122}$ exchange reactions at different temperatures. Providing that the exchange can be effected, variations in the $\text{Te}^{130}/\text{Te}^{122}$ ratio in equilibrium isotope reactions of up to 2.6% at 20°C are predicted. Calculations are also made of kinetic isotope effects predicting values as high as 1.0175 for the ratio of isotopic rate constants.

THE

PROCEEDINGS OF THE

ANNUAL MEETING OF THE

AMERICAN ASSOCIATION OF

PHYSIOLOGISTS

Held at the

Hotel Hamilton, New York City,

December 29, 1900.

Published by the

Association of American Physicians.

1901.

Entered as Second-Class Matter, May 1, 1896.

Postage paid at New York, N. Y., and at additional mailing offices.

Acceptance for mailing at special rate of postage provided for in

Postoffice Department Circular No. 1109, approved October 3, 1900.

Postmaster: This publication is entered as Second-Class Matter.

under Postoffice Department Circular No. 1109, approved October 3, 1900.

and is accepted for mailing at special rate of postage provided for in

Postoffice Department Circular No. 1109, approved October 3, 1900.

under Postoffice Department Circular No. 1109, approved October 3, 1900.

and is accepted for mailing at special rate of postage provided for in

Postoffice Department Circular No. 1109, approved October 3, 1900.

Experimental verification of some of the theoretical predictions was obtained. In the chemical reduction of potassium tellurite at room temperature, the $\text{Te}^{122}\text{O}_3^-$ ion was found to be reduced 1.0071 ± 0.0003 times faster than the reduction rate of $\text{Te}^{130}\text{O}_3^-$. In a microbiological reduction of tellurite ion, the $\text{Te}^{122}\text{O}_3^-$ ion was found to be reduced 1.0060 ± 0.0005 times faster than the reduction rate of $\text{Te}^{130}\text{O}_3^-$.

On the bases of the theoretical predictions, the effects noted in both reduction studies are related to tellurium oxygen bond breakage.

Of four natural samples, three were identical in their $\text{Te}^{122}/\text{Te}^{130}$ ratio; whereas, the fourth was enriched by 0.4 percent in the lighter isotope.

ACKNOWLEDGEMENTS

The author wishes to express his sincere thanks to his supervisor, Dr. H. R. Krouse, for suggesting this project and for his inspiration, guidance, and assistance in the course of this investigation. The assistance of various members of the Department of Computing Science; the technical services of P. Alexander, glassblower; and assistance of R. McCready of the Department of Microbiology in obtaining bacterial cultures and medium are also greatly appreciated.

Finally, I wish to acknowledge the assistance of my wife, Marilyn, and Mrs. P. Babet in the preparation of the manuscript.

TABLE OF CONTENTS

INTRODUCTION	Page 1
A. Equilibrium Isotope Effects	3
B. Kinetic Isotope Effects	4
C. Mechanisms of Isotope Fractionation in Nature	6
1. Physical processes	6
a. Diffusion	6
b. Isotopic vapour pressures	7
c. Miscellaneous	9
2. Chemical processes	9
a. Equilibrium isotopic exchange	9
b. Kinetic isotope effects	13
3. Biological processes	13
D. The Group VIA Elements	15
1. General chemistry	15
2. Isotope fractionation studies	18
a. Sulphur	18
b. Selenium	26
3. Tellurium	28
a. History and occurrence	28
b. Uses	30
c. Physical properties	30
d. Chemical properties	34
e. Absolute abundances of the isotopes of tellurium	35
E. Advantages of looking at sulphur, selenium, and tellurium isotopes simultaneously in natural deposits	37
THEORY	41
A. Theory of Equilibrium Isotope Effects	42

THE UNIVERSITY OF CHICAGO

PHILOSOPHY DEPARTMENT

1990-1991

PHILOSOPHY 101

Introduction to Philosophy

This course is designed to provide students with a broad overview of the major areas of philosophy, including metaphysics, epistemology, ethics, and political philosophy.

The course will be taught by Professor [Name]

and will meet on [Days]

from [Time] to [Time] in [Room]

Students are encouraged to

attend the course as a prelude to

more advanced study in philosophy.

For more information, please

contact the Philosophy Department

at [Address]

or call [Phone Number]

at [Area Code] [Number]

or

write to [Address]

[City, State, Zip]

Students are also encouraged to

visit

the Philosophy Department's

website at [URL]

for more information.

The University of Chicago is an equal opportunity institution. All persons are encouraged to apply.

For more information, please contact the Philosophy Department at [Address]

B. Evaluation of Partition Function Ratios	45
C. Kinetic Isotope Effects	49
D. Isotope Fractionation During First Order Competing Reactions	56
THEORETICAL CALCULATIONS	59
A. Calculation of Equilibrium Exchange Constants	59
B. Calculation of Ratios of Isotopic Rate Constants	64
EXPERIMENTAL	68
A. Extraction of Tellurium from Natural Samples	69
B. Chemical Reduction of K_2TeO_3	71
C. Reduction of K_2TeO_3 by <u>Scopulariopsis</u> <u>brevicaulis</u>	73
1. Reduction of K_2TeO_3 on liquid medium	75
2. Extraction of elemental tellurium reduced from K_2TeO_3 in the biological investigation	76
D. Preparation of TeF_6	77
E. Mass Spectrometric Analysis	80
RESULTS AND DISCUSSION	85
A. Chemical Reduction of Potassium Tellurite	85
B. Microbiological Reduction Study	88
C. Natural Isotopic Abundance Studies	91
BIBLIOGRAPHY	93
APPENDIX I	
CALCULATIONS OF ISOTOPIC VIBRATIONAL FREQUENCY SHIFTS FOR TELLURIUM - CONTAINING MOLECULES	A1
A. Diatomic Molecules	A1
B. Polyatomic Molecules	A5
TeF_6	A7

TeBr ₄	A14
SCTe	A18
H ₂ Te	A22
TeO ₃ ⁼	A30

APPENDIX II

IBM 7040 COMPUTER PROGRAMS FOR CALCULATION OF ISOTOPIC VIBRATIONAL FREQUENCIES AND PARTITION FUNCTION RATIOS	A36
A. Calculation of Isotopic Vibrational Frequencies and Partition Function Ratios for Diatomic Species	A37
B. Computation of Isotopic Vibrational Frequencies for XY ₆ Molecules	A40
C. Computation of Isotopic Vibrational Frequencies for XY ₄ Molecules	A43
D. Computation of Vibrational Frequencies for XYZ Molecules	A46
E. Computation of Force Constants for Non- linear XY ₂ Molecules	A49
F. Computation of Isotopic Vibrational Frequencies and Partition Function Ratios for Non-linear XY ₂ Molecules	A55
G. Computation of Force Constants, Apex Angle, Isotopic Vibrational Frequencies, and Partition Function Ratios for Pyramidal XY ₃ Molecules	A58
H. Computation of Partition Function Ratios for General Molecules	A63



LIST OF TABLES

	Page
I. Occurrence of Tellurium	31
II. Absolute Abundances of the Isotopes of Tellurium	38
III. Q_2'/Q_1' for Te-containing Molecules	61
IV. Equilibrium Constants for $\text{Te}^{130} \xrightleftharpoons{\quad} \text{Te}^{122}$ Exchange Reactions	62
V. Minimum Values of k_{130}/k_{122}	65
VI. Summary of the Calculated Ratios of Rate Constants for the Reduction of K_2TeO_3	67
VII. Ion Species Resulting from the Ionization of TeF_6	84
VIII. $\text{Te}^{130}/\text{Te}^{122}$ Fractionation Study	86
IX. Calculation of Vibrational Frequency Shifts for Te^{130} and Te^{122} -containing Molecules	A6

LIST OF FIGURES

	Page
Figure 1. O^{18} Growth Rings in the Jurassic Belemnite	12
Figure 2. The Distribution of S^{34} in Nature	20
Figure 3. The Biological Sulphur Cycle	22
Figure 4. Variations in the Se^{82}/Se^{76} Ratio Found For Natural Samples	27
Figure 5. Fluorine Line	78
Figure 6. Typical Tellurium Spectra	82
Figure 7. Te^{122}/Te^{130} Fractionation Study	87
Figure 8. Variation of Isotopic Composition of Product with Percent Reaction	89
Figure 9. Normal Vibrations of an Octahedral XY_6 Molecule	A 8
Figure 10. Normal Vibrations of a Tetrahedral XY_4 Molecule	A15
Figure 11. Relationship Between Force Constants for H_2Te	A26
Figure 12. Normal Vibrations of a Pyramidal XY_3 Molecule	A32

INTRODUCTION

The concept of isotopes, new and revolutionary in 1910, is today basic to our understanding of the structure of matter.

The first direct evidence of the existence of isotopes came from studies of radioactive elements and their decay products. Soon the chemists investigating the decay products found pairs of radioactive elements which were inseparable and appeared to be identical in their chemical properties. Soddy (100) in 1910 suggested that such substances differing in their physical properties but having the same chemical properties be called "isotopes". He suggested that the existence of isotopes was a general phenomenon and that we could also expect the stable elements to be mixtures of these inseparable species, or isotopes.

J. J. Thomson in 1912 (128) using his positive ray apparatus, showed that neon was indeed a mixture of at least two kinds of atoms having atomic weights of 20 and 22 since the positive ion beam formed from neon was resolved into two beams. This was the first evidence for the existence of isotopes among the stable elements.

Aston (2) and Dempster (25), subsequently examined the isotopes of many other elements. During the two decades to follow, practically all the elements had been investigated. Today, the known stable elements consist of mixtures of from one to ten isotopes and these total to about 300 different stable isotopes in addition to the radioactive species. The radioactive disintegration studies suggested that isotopic abundances of an element in a deposit could be altered by radioactive decay. Richards (90) found that lead associated with thorium ores had a higher atomic weight than that associated with uranium ores. This was reasonable since Th^{232} decay terminates in Pb^{208} ; whereas U^{238} decays finally to Pb^{206} . By knowing the amount of radioactively produced lead present relative to the amount of undecayed parent, the age of lead deposits can be determined.

Similarly, Aldrich and Nier (1) observed that the amount of Ar^{40} associated with K^{40} was up to three times the concentration of Ar^{40} in atmospheric argon. The amount of Ar^{40} produced by the radioactive decay of K^{40} has been used to date the time of formation of potassium minerals. Many other radioactive isotopes have been used for dating purposes.

However, despite all the information about isotopes that had been discovered prior to 1930, it was believed that isotopes were identical in their chemical properties and that the stable isotopes of the elements were of constant abundance in nature. But soon after the discovery of deuterium, the heavy isotope of hydrogen, by Urey, Brickwedde, and Murphy (112) in 1932, it was shown by theory and experiment that the isotopes of the light elements did differ in their chemical properties and that the separation of these isotopes by chemical means was possible.

A. Equilibrium Isotope Effects

In 1933 Urey and Rittenberg (114) calculated the equilibrium constants for the reaction



at different temperatures. They found these to differ from unity which indicated that the isotopes of hydrogen were different chemically. The reaction was carried out in the laboratory at various temperatures by Rittenberg, Bleakney, and Urey (91) and the equilibrium constants determined. These were found to be close to those calculated theoretically. This was the beginning of many

isotope fractionation investigations which compared experimental results with theoretical calculations.

Urey and Greiff (116) calculated the equilibrium constants for some of the light elements such as lithium, boron, carbon, nitrogen, chlorine, and bromine. Their results indicated equilibrium fractionations varying from 1% to 10% and experimental confirmations of some of these were soon obtained. For example, Weber et al (127) found CO_2 in equilibrium with water to be enriched in O^{18} by 4.6% at 0°C as compared with 4.4% predicted by theory. Other equilibrium isotope effects and data are summarized by Urey (118). Many isotope exchange reactions involving the isotopes of the light elements have been studied in recent years and the agreement between theory and experiment has been excellent. To date, (with the exception of this thesis) selenium is the heaviest element for which exchange reactions have been investigated (60).

B. Kinetic Isotope Effects

In 1932 when Urey and Washburn (113) were doing their experiments with deuterium and protium, they discovered that partially electrolyzed water was enriched in deuterium, protium having been evolved faster than

deuterium at the cathode. Two years later, Bach, Bonhoeffer, and Fajans (5) showed that H_2 reacted over three times faster than D_2 with bromine. The same year Farkas and Farkas (37) reported similar results in the photochemical reactions of H_2 and D_2 with chlorine. Urey and Teal (117) and Eidinoff (31) have reviewed the extensive work done on the relative reaction rates of H_2 and D_2 - containing compounds.

Kinetic isotope effects were similarly found for other lighter elements. In 1949 Lindsay, McElheran, and Thode (65) and Bigeleisen and Friedman (12) reported carbon isotope effects in the decomposition of oxalic acid and the decarboxylation of malonic acid. These results were reported on further by Lindsay, Bourns, and Thode (63,64) in 1951 and 1952 and were in excellent agreement with theory.

To date, much literature is available on the mathematical calculation of isotopic rate constants. In 1935 Eyring (36) set forth the theory of absolute rates from which the ratio of rate constants for reactions of isotopic molecules can be calculated, and this has been revised by Bigeleisen (11) to an expression which is more convenient to use. Several authors (23, 92, 106) have written reviews on kinetic isotope effects.

C. Mechanisms of Isotope Fractionation in Nature

Natural isotope fractionation mechanisms can be subdivided into physical, chemical, and biological processes.

1. Physical processes of natural isotope fractionation

a. Diffusion: The mass dependence of diffusion in the solid, liquid, or gaseous state can lead to isotopic fractionation and this is one of the principles used in commercial isotope separation. Studies of natural diffusion phenomena for many isotopes have been reported. An interesting example is the studies of nitrogen isotopes by Hoering (53). He discovered that nitrogen contaminants in natural gas were enriched in N^{14} over the nitrogen found in the associated crude oil. He suggested that migration of the oil and gas through porous deposits was responsible. Migration by molecular flow in which the mean free path of the gas is greater than the pore dimensions results in flow rates of the isotopic molecules which are inversely proportional to the square root of the masses. This postulate was later verified by a laboratory diffusion experiment and he was able to produce to some degree the fractionation found in nature.

With lighter isotopes and low temperature conditions, the isotopic alteration can be quite high. This is exemplified by a recent lithium diffusion experiment in titanium dioxide by Johnson and Krouse (55). Lithium was allowed to diffuse to a depth of 1.7 mm at room temperature and the innermost lithium was found to be 7 percent enriched in Li^6 in comparison to the undiffused lithium showing that the Li^6 diffuses quite noticeably faster than the Li^7 .

b. Isotopic vapour pressures: Friedman (39) suggested that the different vapour pressures of the isotopic compounds might influence the natural abundance of each isotope. He showed that there was a gradual depletion of the deuterium content in rainfall samples progressing from the Pacific Coast of North America inland over the Rocky Mountains. He found further that ocean waters near the equator were found to be enriched in deuterium in comparison to ocean waters polewards. Both these phenomena are caused by the preferential evaporation of H_2O over HDO . In the rainfall study, the HDO condenses faster enriching the coastal regions in HDO while the remaining vapour enriched in H_2O moves further inland.

The oxygen isotopes of the earth's water bodies behave similarly. Epstein (34) reported that the O^{18} content of the ice is very low in the Arctic regions relative to ocean waters and that the actual value in the ice cap varies in a cyclic fashion with depth. This cyclic phenomenon which is due to seasonal changes may be explained by thinking of the atmosphere between the equator and the poles as a distillation column. Water is evaporated from the oceans at the equator, rises upward at the equator, and spreads out toward the poles. In the distillation of the water, the simple process separation factor α is given by the ratio of vapour pressures of the two isotopic water species as

$$\alpha = \frac{P_{H_2O^{16}}}{P_{H_2O^{18}}} = \begin{matrix} 1.008 \text{ at } 25^\circ\text{C} \\ 1.011 \text{ at } 0^\circ\text{C} \end{matrix}$$

Since H_2O^{16} has the higher vapour pressure, the O^{18} is favoured in the liquid and will be depleted in the vapour. As rain falls in the different latitudes, the vapour spreading out towards the poles becomes more and more depleted in the heavy isotope. We therefore have a multi-stage process. The extent of the isotope fractionation which in this case is the depletion of O^{18} in snow fall-


ing on the Greenland ice caps will depend on the number of stages in the process and on the difference in temperature between the equator and the poles. Thus, the depletion of O^{18} in the snow and ice in the Arctic will, in general, be higher in winter than summer. Seasonal fluctuations in the O^{18} content with depths have shown up in the layers of ice in the Greenland glacier. From such studies, estimates of the age of glaciers can be determined by simply counting O^{18} concentration rings. Such fractionations caused by differences in vapour pressure are termed "Rayleigh" fractionations.

c. Miscellaneous: It is expected that differences in the isotopic solubilities, melting and sublimation points might be other mechanisms of natural isotope fractionation. Very little work to date has been reported on these physical processes.

2. Chemical processes


a. Equilibrium isotopic exchange: Friedman suggested further that equilibrium isotope exchange reactions would produce fractionation of isotopes in natural processes. For example, a study of fumaroles in Yellowstone Park showed the deuterium content to be depleted by as much as 45%. He suggested that this might

be the result of the equilibrium exchange reaction



The equilibrium constant for exchange reactions is dependent upon temperature, and if this hypothesis is the only contributing factor, theoretical considerations, based on the observed deuterium depletions, place the temperatures of these fumaroles above 400°C.

In the exchange of the oxygen isotopes between carbonate and water, the heavy isotope O^{18} is favored in the carbonate over that of that water. This reaction can be written in the form



The equilibrium constant has been determined by Clayton (20) to be 1.0301 at 25°C. meaning that at 25°C. limestone and calcium carbonate material will be enriched in O^{18} over that of sea water by about 3%. The exact extent of this enrichment will depend on the temperature of the water at the time of deposition. The reaction has been studied in detail by Urey (119) and he suggested that it might be used in the determination of temperatures and that the accurate measurement of the O^{18} content of

The following is a list of the names of the persons who have been elected to the office of

President

of the Association for the year 1900.

The names of the persons who have been elected to the office of

President are as follows:

1. Mr. J. H. Smith

2. Mr. J. H. Smith

3. Mr. J. H. Smith

4. Mr. J. H. Smith

5. Mr. J. H. Smith

6. Mr. J. H. Smith

7. Mr. J. H. Smith

8. Mr. J. H. Smith

9. Mr. J. H. Smith

10. Mr. J. H. Smith

11. Mr. J. H. Smith

12. Mr. J. H. Smith

13. Mr. J. H. Smith

14. Mr. J. H. Smith

15. Mr. J. H. Smith

suitable marine carbonate sediments would indicate the temperature of the seas at the time of deposition.

According to Clayton the temperature T is given by the equation:

$$\ln K = 2725 T^{-2}$$

As a direct result of this temperature dependence, the concept of the O^{18} or paleo-temperature scale was formulated.

Studies were carried out by Urey et al (120), Epstein et al (32), and McCrea (73) using well preserved samples of belemnites from the U. S., Sweden, Denmark, England, France, Holland, and Germany and these showed that the temperatures which were determined seemed reasonable.

A fascinating study was made by Epstein (33) of the O^{18} content in the carbonate shells generated by marine animals. The subject of investigation was a Jurassic belemnite, an extinct animal which flourished about 120 million years ago. Successive growth rings of the shells showed a cyclic variation in the heavy oxygen (O^{18}) content which were associated with seasonal variations in temperatures. Figure 1 shows these seasonal vibrations in temperature as indicated by the O^{18} content of the




Fig. 1.- 018 growth rings in the Jurassic belemnite.

Taken from Thode (108)

growth rings of the belemnite. Such an oxygen isotope study, therefore, not only gives us information about climatic conditions in various parts of the world millions of years ago, but tells us something about the lives and habits of animals that now are extinct.

b. Kinetic isotope effects: Due to the difficulty in observing purely chemical kinetic isotope effects in nature, very little has been reported in the literature. Numerous reports have been given for kinetic isotope fractionation in biochemical processes and will be discussed later in connection with sulphur isotope fractionation.

3. Biological processes

Biological and microbiological isotope fractionations are perhaps due to both physical and chemical processes. Because of the complexity of the mechanisms, there is insufficient data to verify whether isotopes are fractionated in such processes as enzyme complexing and diffusion through biological membranes. Significant isotope fractionations have been realized with carbon, nitrogen, oxygen, and sulphur in biochemical processes and the results in many instances are consistent with similar inorganic chemical processes.

The first of these is the fact that the
theoretical model is based on the assumption
that the system is in a steady state. This
assumption is not valid in the case of a
transient process. The second is the fact
that the model is based on the assumption
that the system is linear. This assumption
is not valid in the case of a non-linear
system.

The third is the fact that the model
is based on the assumption that the
system is time invariant. This assumption
is not valid in the case of a time
variant system. The fourth is the fact
that the model is based on the assumption
that the system is causal. This assumption
is not valid in the case of a non-causal
system. The fifth is the fact that the
model is based on the assumption that
the system is stable. This assumption
is not valid in the case of an unstable
system.

The sixth is the fact that the model
is based on the assumption that the
system is deterministic. This assumption
is not valid in the case of a stochastic
system. The seventh is the fact that
the model is based on the assumption
that the system is continuous. This
assumption is not valid in the case of
a discrete system. The eighth is the
fact that the model is based on the
assumption that the system is single
input single output. This assumption
is not valid in the case of a multi
input multi output system. The ninth
is the fact that the model is based on
the assumption that the system is
linear time invariant. This assumption
is not valid in the case of a non-linear
time variant system. The tenth is the
fact that the model is based on the
assumption that the system is stable.
This assumption is not valid in the
case of an unstable system.


Nier and Gulbransen (83) and two years later, Nier and Murphy (84) found that in general, limestones were enriched and plants depleted in the heavier carbon isotope. They reported variations up to 5% in the C^{12}/C^{13} abundances. Many workers have since re-affirmed these results and can be found in the literature (6,22,38,62,71,80,86,97,107,110,130,133,134,135). One of the many interesting results of these investigations was that both land and marine plants tend to concentrate C^{12} , although the latter do this to a lesser degree. This phenomenon may be attributed to differences in the cellular structures of the two types of plants, assuming that the concentration of the C^{12} is due to kinetic and/or equilibrium isotope effects in the photosynthesis of atmospheric CO_2 .

Much investigation of the isotope fractionation due to bacteria has been carried out. The most fruitful of these investigations have been carried out with sulphur, and this will be discussed in more detail later.

D. The Group VIA Elements

1. General chemistry (47), (96)

The group VI A elements include oxygen, sulphur, selenium, tellurium, and polonium and are characterized by 6 valence electrons. They differ in nuclear charge, in size, and in the number of inner electron "shells". Little is known about polonium since work on it has been retarded because of its high radioactivity. It occurs naturally in uranium minerals such as pitchblende but only to the extent of 5×10^{-9} per cent of the mineral. Thus its occurrence is so small that its significance in stable isotope fractionation processes is negligible.



The Group VIA Elements

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
JANUARY 1950
RECEIVED
FROM THE
LIBRARY OF THE
UNIVERSITY OF CHICAGO
LIBRARY



The firmness with which the outer electrons are held increases with increasing nuclear charge, but is diminished as the distance from the nucleus and the number of inner electron shells increases. Since the two latter effects predominate, it may be predicted that the electrons are most tightly held by the oxygen atom and most easily lost (or shared) by the tellurium atom. This assumption accounts for and is in accord with the systematic variation in physical and chemical properties which these elements show.

The elements of the group readily form compounds or ions by sharing or adding 2 electrons, provided by atoms of other elements, to complete the octet of the inert-gas structure thereby acquiring an oxidation state of -2. This tendency is most prominent in the case of the smallest atom, that of oxygen, and least in the case of the largest atom, that of tellurium. In addition, the atoms of sulphur, selenium, and tellurium may lose (or share) their valence electrons so as to acquire oxidation states up to +6. Oxygen does not show this tendency, because its electrons are so tightly bound that no other element, with the possible exception of fluorine, is able to remove them. The change in oxidation state

of an element during a chemical reaction is one of the factors which determines the extent to which isotope abundances may be altered in the laboratory or by natural processes.

All of the elements of the group show allotropy. Just as oxygen can exist as diatomic O_2 and triatomic O_3 , the other elements can be obtained in more than one form, the forms differing either in the number of atoms per molecule, or in the arrangement of molecules in the solid. There is, going from sulphur to tellurium, an increasing disposition toward formation of long chains of atoms held together by covalent bonds. In general, the larger the atom, the less the tendency to form multiple bonds and the greater the tendency of each atom to be bound to more than one atom. Oxygen stands alone from the group in being a diatomic gas at room temperature.

As would be expected because of the increasing number of electronic shells, the ionic radii increases from $1.40\text{\AA}$ for oxygen, $1.84\text{\AA}$ for sulphur, $1.98\text{\AA}$ for selenium, to $2.4\text{\AA}$ for tellurium (47). Also of interest, the oxidation potential for each of these elements is as follows: oxygen, -1.23 volts; sulphur, -0.14 volts; selenium, +0.40 volts; and tellurium, +0.72 volts (96).

...the ... of ... in ...

...the ... of ... in ...

...the ... of ... in ...

...the ... of ... in ...

...the ... of ... in ...

...the ... of ... in ...

...the ... of ... in ...

...the ... of ... in ...

Thus, there is a much greater tendency for oxygen to form H_2O than for tellurium to form H_2Te . Unlike H_2O and H_2S , H_2Se and H_2Te are better reducing agents than hydrogen.

From the foregoing discussion, it is readily seen why the properties of sulphur, selenium, and tellurium vary from the decidedly non-metallic nature of sulphur to the mildly metallic character of tellurium. For example, the free element sulphur is a non-conductor of electricity, selenium is a poor conductor (but its conductivity increases proportionately to its illumination), and tellurium is somewhat better than selenium.

2. Isotope fractionation studies

In considering isotope fractionation in the chemical group VIA, oxygen has received extensive study and some examples have been previously cited. It would seem, however, that tellurium isotopes should resemble sulphur and selenium in their behavior because of the similarities of their valence states. Therefore, some of the significant findings of sulphur and selenium isotope fractionation studies are reviewed here.

a. Sulphur: Extensive studies of the variations in the isotopic abundances of sulphur have been

...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...

...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...


...the ... of ... and ...

...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...

...the ... of ... and ...
...the ... of ... and ...
...the ... of ... and ...

carried out. It was clear that differences in the chemical properties of the sulphur isotopes did exist and that isotope fractionation could be expected in natural processes.

For example, calculations of the isotope exchange constant for the reaction



gives K at 25°C to be 1.071, thus favoring the enrichment of S³⁴ in the sulphate by about seven percent (111).

It was therefore suspected that a trend toward this favoured distribution existed in nature with sulphates enriched and sulphides depleted in the heavy isotope of sulphur, S³⁴.

The initial data published by Thode, Macnamara, and Collins (104) showed that the S³²/S³⁴ ratio in terrestrial samples varied by as much as 5%. As predicted, sulphates were generally found to be enriched in S³⁴ while sulphides were depleted in the heavier isotope. Figure 2 shows the general distribution of sulphur isotopes in nature. The sulphur isotope distribution data are expressed in terms of δS³⁴ ‰ defined as follows:

THE UNIVERSITY OF CHICAGO

DEPARTMENT OF THE HISTORY OF ARTS

RECEIVED

1964

1964

THE UNIVERSITY OF CHICAGO

DEPARTMENT OF THE HISTORY OF ARTS

RECEIVED

1964

1964

1964

1964

1964

1964

1964

1964

1964

1964

1964

1964

1964




FIG. 2 THE DISTRIBUTION OF S^{34} IN NATURE
Taken from Thode (108)

$$\delta S^{34} \text{ ‰} = \frac{(S^{34}/S^{32})_{\text{sample}} - (S^{34}/S^{32})_{\text{standard}}}{(S^{34}/S^{32})_{\text{standard}}}$$

Positive and negative δ -values mean that the samples are enriched or depleted in S^{34} respectively with respect to the standard.

It was reported by Macnamara and Thode (68) that the sulphur in meteorites had a remarkably constant S^{32}/S^{34} ratio which was close to the average S^{32}/S^{34} ratio for terrestrial samples. These facts led them to believe that the meteoritic S^{32}/S^{34} ratio represented the primordial abundance of terrestrial sulphur before any fractionation occurred or the ratio of the sulphur isotopes in the earth at the time that it was formed. It is clear that since the earth was formed, the sulphur in the crust has been fractionated in biological and geological processes due to differences in the chemical properties of the isotopes.

The exchange cited does not take place under normal conditions (124) and all efforts to establish it under laboratory conditions for a reasonable length of time failed. Therefore, Tudge and Thode (111) suggested that the following biological sulphur cycle might well provide the mechanism for this fractionation in nature (Fig. 3).




FIG. 3 THE BIOLOGICAL SULPHUR CYCLE IN NATURE
Taken from Thode (108)

According to Thode (108) the bacterial reduction process is found to be a unidirectional, not an equilibrium process, and isotope fractionation occurs because the light-isotope containing compounds react faster than the species containing the heavier isotope. Thode goes on to say that in particular, the bacterial reduction of sulphate is largely responsible for the isotopic variations which occur in the oceans, evaporites, and materials in the sediments.

Many studies have been made of isotope fractionation in the sulphur cycle. Anaerobic reducing and photosynthetic hydrogen sulphide oxidizing bacteria abound in the Cyrenacian lakes of Africa. These produce a sedimentary layer of elemental sulphur (15) and analysis of samples from these lakes gave direct evidence for the fractionation of sulphur in the sulphur cycle. Macnamara and Thode (69) found the S^{34} content of the elemental sulphur to be 3.2% less than that of the soluble sulphate from the same lake.

Further evidence is provided in the salt domes of Louisiana and Texas (107) where elemental sulphur was depleted by about 4% and sulphide by about 5% in S^{34} relative to the associated gypsum. Two possibilities

exist for the formation of this sulphur by reduction of the sulphate: - either the high temperature reduction of the sulphate by organic matter (oil fields associated with the salt domes), or by the bacterial reduction of the sulphate with petroleum providing a source of carbon and free energy for life. The first of these mechanisms, however, seems incapable of causing the large fractionation observed; whereas, the latter is quite capable of doing so, as shown by the Cyrenacian Lake study. Miller (78) added further evidence to the "bacterial reduction" postulate by isolating anaerobic sulphate-reducing bacteria in these deposits. Thus the production of carbon dioxide by these sulphate-reducing bacteria alters the calcium sulphate to a porous calcitic limestone with elemental sulphur embedded in it and this accounts for the type of deposit that is present in these salt domes today.

The hypothesis that sulphate-reducing bacteria were responsible for the sulphur isotope fractionation has been substantiated by laboratory experiments. Thode, Kleerekoper, and McElheran (105) showed that Desulphovibrio desulfuricans while reducing sulphate, produced H_2S which was 1% richer in S^{32} than the nutrient. In subsequent studies, Harrison and Thode (44) found that this fraction-

ation varied with reaction conditions and that the maximum enrichment was 2.5% at 25°C. Kaplan and Rittenberg (57) in 1964 reported enrichments as high as 4.6%. They showed also that other micro-organisms such as S. cerevisiae were capable of carrying out fractionation of oxidized sulphur compounds in reduction experiments with equal or greater efficiency.

McElheran in 1951 (74) was unable to find any fractionation in the oxidation of sulphur to sulphate by Thiobacillus but in 1964 Kaplan and Rittenberg (57) reported an enrichment of 1.8% of S^{32} in oxidizing H_2S to $SO_4^{=}$ by chemosynthetic oxidation and an enrichment of 1.0% in photosynthetic oxidation. Nakai and Jensen (96) published results in 1964 for the bacterial reduction and oxidation of sulphur using marine mud cultures. Values for the ratio of the isotopic (S^{32} and S^{34}) rate constants of about 1.020 were found for the reduction experiments, and values ranged from 1.0003 to 1.0017 for the oxidation experiment. Fractionation factors between sulphate and sulphide in bottom sediments vary between 1.03 and 1.04 for sea samples and 1.005 to 1.009 for fresh water samples, indicating that the sulphate produced becomes more enriched in S^{32} than does sulphur or sulphide.

b. Selenium: An investigation into the isotopic fractionation of selenium was conducted by Krouse and Thode (60). They reported $\text{Se}^{82}/\text{Se}^{76}$ ratios as found in figure 4. The $\text{Se}^{82}/\text{Se}^{76}$ ratio varied 1.5% in the samples examined. The distribution bears some similarity to that of the sulphur isotopes. Samples of plant materials and soil showed the largest variation in isotope ratios; whereas, selenides from massive sulphide ores showed little or no variation with respect to meteoritic selenium.

They further reported that Se^{76}O_3 was reduced 1.5% faster than Se^{82}O_3 to elemental selenium. This result is in agreement with a simple theoretical calculation. This compares to a 2.3% kinetic isotope effect in the chemical reduction of S^{32}O_4 and S^{34}O_4 reported by Harrison and Thode (44).

Further investigation into the isotope fractionation of selenium has been carried out by Rees (89). Among other experiments, he carried out investigations into the kinetic isotope effects in the reduction of Se^{IV} to Se^0 , and into the isotope exchange reaction between Se^{IV} and Se^{VI} . He reported a kinetic isotope effect varying from 1.6% at 40°C to 2.0% at 24°C for a 20% reduction of sodium selenite,




FIG. 4 VARIATIONS IN THE $^{82}\text{S}_0 / ^{80}\text{S}_0$ RATIO FOUND FOR NATURAL SAMPLES

and phase fractionations of 2.7% and 4.2% respectively for 59 - and 146 - day equilibrations of Se^{VI} and Se^{IV} . In each case, the higher valence states were enriched in the heavier Se^{82} .

Because of the numerous reports of isotopic fractionation of sulphur and selenium, a similar fractionation of tellurium isotopes by natural processes is strongly predicted.

3. Tellurium

a. History and occurrence - The discovery of tellurium was made late in the eighteenth century (76). A peculiar mineral known as white gold-ore or grey gold-ore occurring in the sandstone near Zalathna, Transylvania, was considered by chemists and mineralogists as a kind of alloy of antimony and bismuth. Various forms were given a host of names but not until 1782 that F.J. Müller von Reichenstein showed that there was neither any bismuth nor antimony in the new mineral, and suspected that a new element had been discovered. Then sometime after 1789, M. H. Klaproth extracted the unknown metal and observed some of its properties from which concluded that the mineral contained a peculiar and distinct metal essentially different from every other metallic substance

hitherto known. It was he who named it tellurium after the earth (Latin - tellus).

Tellurium occurs in only a few places and in small quantities; it is less abundantly distributed than any other element in the sulphur family. According to the data of spectrographic analysis, it is absent in the spectra of the sun and the stars, however, it could be explained by the low sensitivity of the spectrographic method with respect to these elements (98). According to F. W. Clarke and H. S. Washington (19), the igneous rocks on the earth's crust contain 8×10^{-2} percent sulphur, $n \times 10^{-8}$ percent selenium, and $n \times 10^{-9}$ percent tellurium; or tellurium is about as abundant as gold. It occurs combined with gold, silver, bismuth, and many other metals. The chief tellurium minerals are the tellurides, but some tellurites and tellurates exist in natural form. These are listed in Table I. Tellurium has not been detected in soils (123). It is found in many volcanic areas and has been found in "minute" amounts in some meteorites (29). It is obtained principally as a by-product of the refining of other minerals with which it combines.

b. Uses - To cite some uses, it is used in coloring glass, preparation of organic dye-stuffs, manufacture of high resistance alloys, staining silver, as a delicate test of sterilization in bacteriology, identification of presence of diphtheria and other germs, in motor fuels as an anti-knock constituent, insecticides, germicides, fungicides and wood preservatives, and in the manufacture of extremely hard alloys having very great tensile strength. Its main use in recent years is in the semiconductor field where it has found use as an intrinsic photoconductive detector (77) and also in rectifiers (96).

c. Physical properties - Tellurium exists primarily as "metallic" tellurium which is the most stable allotrope. This form is grey black in color with a metallic luster and of hexagonal crystal structure. In 1913 Cohen and Kröner (21) claimed that tellurium existed in two dynamic allotropes, and according to Mellor (76) tellurium is obtained in the amorphous form as a black powder when tellurium is precipitated from an acid solution. The amorphous form suffers no change upon heating until it is melted (m.p. 452°C) and then it is transformed into the very brittle and easily powdered white-grey crystalline form.

Table I OCCURRENCE OF TELLURIUM (76), (98)

<u>Native Element</u>	Te
<u>Tellurides</u>	
Altaite	PbTe
Arseniotellurite	$\text{Te}_2\text{As}_2\text{O}_7$
Arsenotellurite	$\text{Te}_2\text{As}_2\text{S}_7$
Calaverite	AuTe_2
Coloradoite	HgTe
Csiklovaite	Bi_2TeS_2
Frohbergite	FeTe_2
Goldfieldite	$\text{Cu}_6\text{Sb}_2(\text{S}, \text{Te})_3$
Goldschmidtite	Au_2AgTe_6
Grünlingite	$\text{Bi}_4\text{S}_3\text{Te}$
Hedleyite	Bi_7Te_3
Hessite	Ag_2Te
Hydrogen telluride	H_2Te
Joséite	Bi_3TeS
Krennerite	$(\text{Au}, \text{Ag})\text{Te}$
Melonite	Ni_2Te_3
Montbrayite	Au_2Te_3
Muthmannite	$(\text{Ag}, \text{Au})\text{Te}$
Nagyagite	$\text{Au}_2\text{Sb}_2\text{Pb}_{10}\text{Te}_6\text{S}_{15}$

Oruetite	$\text{Bi}_3\text{Te}_3\text{Bi}_2\text{S}_3$
Petzite	$(\text{Ag}, \text{Au})_2\text{Te}$
Pilasonite	Bi_3Te_2
Rickardite	Cu_4Te_3
" Stutzite	Ag_4Te
Sylvanite	AuAgTe_4
Talapite	$\text{Bi}(\text{S}, \text{Te})_3\text{Ag}_3$
Telluro bismuthite	Bi_2Te_3
Tetradymite	Bi_2Te_2
Vondiestite	$(\text{Ag}, \text{Au})_5\text{BiTe}_4$
Wehrlite	$\text{Bi}_7\text{Te}_7\text{Ag}$
Weissite	Cu_2Te

Mixtures

Coolgardite	coloradoite, petzite, calaverite, sylvanite
Henryite	lead telluride and pyrites
Kalgoorlite	coloradoite, petzite, and tellurium
White tellurium (Gelberg, mullerite)	antimonial telluride of gold, silver, and lead

Tellurites

Durdenite	$\text{Fe}_2(\text{TeO}_3)_3 \cdot 4\text{H}_2\text{O}$
Tellurite	TeO_2


1. The first part of the paper discusses the importance of understanding the underlying mechanisms of the observed phenomena. This is followed by a detailed analysis of the data, which shows a clear trend towards a more integrated approach. The results of the study are presented in the following section, highlighting the key findings and their implications for future research. Finally, the paper concludes with a summary of the main points and a call to action for the research community.

2. The second part of the paper focuses on the methodological aspects of the study. It describes the data collection process, the statistical models used, and the validation procedures. The authors emphasize the robustness of the findings by comparing the results with previous studies and conducting sensitivity analyses. The paper also discusses the limitations of the study and suggests directions for future work. The overall structure of the paper is designed to provide a comprehensive overview of the research, from the initial motivation to the final conclusions.


Dunhamite



Emmonsite




Mackayite

Tellurates


Ferrotellurite



Magnolite



Montanite



1898

1899

1900

1901

1902

1903

1904

1898

1899

1900

1901

1902

1903

1904

Textbooks as recently as 1953 claimed a hexagonal lattice and the existence of two polymorphic modifications α and β with a transition point at 354°C . However, Bose (14) in 1941 showed by x-ray diffraction that precipitated tellurium is not amorphous but is hexagonal in a finely divided state, ie. it is the same as metallic tellurium, and Dodwell (27) proved in 1945 that tellurium does not change its structure when heated up to 360°C and consequently does not have any polymorphic varieties. Crystallization of tellurium fused in sodium hydrosulphate and cooled in carbon dioxide results in the formation of a crystalline star resembling that of antimony.

d. Chemical properties - In agreement with the general observation that the basic properties of the natural families of elements increases with atomic weight, tellurium exhibits a greater basicity than selenium and sulphur.

H. Davy (24) was the first to discover hydrogen telluride (H_2Te), which he formed by the electrolysis of water plus acid using a tellurium cathode. H_2Te is also formed by the action of acids on tellurides, but the electrolytic process gives the best yields. It is a colorless gas with a foul smell resembling that of hydrogen sulphide.

Heating of tellurium in the presence of oxygen produces tellurium dioxide, TeO_2 . It forms small, colorless, octahedral crystals. The hydrated form of tellurium dioxide is tellurous acid, H_2TeO_3 . Tellurium monoxide is formed when tellurium sulphite is heated between 180° and 230°C in vacuo ($\text{TeSO}_3 \longrightarrow \text{TeO} + \text{SO}_2$). It is a brown, porous mass, stable in dry air but decomposes when heated into tellurium and tellurium dioxide. Tellurium trioxide (TeO_3) is formed when telluric acid (H_2TeO_4) is heated above 360°C but below red heat. If the temperature be too high, the trioxide decomposes into the dioxide. TeO_3 is orange-yellow and crystalline in form.

Tellurium combines with the halogens to produce tellurium halides. Of particular interest in this work is the relatively unreactive tellurium hexafluoride (TeF_6) gas and its possible use in mass spectrometric determination of the tellurium isotopes. Both hexa-halides and tetra-halides are possible. In the case of fluorine, TeF_6 is formed at temperatures reaching 200°C in excess fluorine, whereas, TeF_4 is formed at temperatures below 180°C in excess tellurium (56).

e. Absolute abundances of the isotopes of tellurium - The first report of tellurium isotopes came in 1925 when Aston (3) discovered Te^{126} ,

Te^{128} , and Te^{130} using his mass spectrograph. As to their relative abundance, he only reported that the latter two were of equal intensity and double that of the first. Then in 1931 he reported Te^{125} (4). Bainbridge (7) in 1932 reported isotopes 122, 123, 124, and 127 and Bartlett (8) in 1934 predicted from $\Delta m/\Delta z$ plots that there should be isotopes Te^{118} and Te^{120} . Dempster (26) in 1936 confirmed Bainbridge's and Bartlett's results using his mass spectrometer but that there was no Te^{118} nor Te^{127} . It was not until 1942 when R. Migeotte (77) using a Bainbridge mass spectrograph reported the eight isotopes with percentage abundances. Four years later, Williams and Yuster (132) confirmed these eight isotopes giving relative abundances. White and Cameron two years later reported similar results (131). The latest values of the isotopes and their percentage abundances were given by Hollander et al in 1953 (54). The results of these investigations are summarized in Table II.

As regarding the atomic mass of each of the isotopes, Duckworth et al (28) in 1951 first reported the masses Te^{126} , Te^{128} , and Te^{130} as 125.9427, 127.9471, and 129.9467 respectively and one year later, Halsted (43) gave the atomic masses for all eight isotopes. Up to 1960, these were the accepted values and no new values were found in

the literature. These values are as given below:

<u>Mass Number</u>	<u>Atomic Mass (amu)</u>
120	119.94288
122	121.94193
123	122.94368
124	123.94278
125	124.94460
126	125.94420
128	127.94649
130	129.94853

E. Advantages of looking at sulphur, selenium, and tellurium isotopes simultaneously in natural deposits

One of the reasons for considering tellurium for an isotope fractionation study is to compare its behaviour with results already obtained with sulphur and selenium. As it has already been shown, a considerable number of studies of the S^{32}/S^{34} variations have yielded much information about natural processes. The preliminary study which has been conducted with Se^{76}/Se^{82} also looks promising in that it shows some similarities to the sulphur isotope distribution.

Whereas sulphur exists in nature from the -2 to the +6 valence states, it is very rare that selenium exists in valence states higher than +4, and tellurium rarely exists

Table II

Absolute Abundances of the Isotopes of Tellurium

Worker Year Reference	Aston 1925 (3)	Aston 1931 (4)	Bainbridge 1932 (7)	Bartlett 1934 (8)	Dempster 1936 (26)
Instrument	Mass spectrograph, photometric detection	Mass spectrograph, photometric detection	Mass spectroscope, magnetic scan- ning, photomet- ric detection	Predicted from $\Delta m/\Delta z$ plots.	Mass spectrometer, electronic detection
Ion source	Te vapour	Te vapour	Te vapour		Spark between tellurium and palladium elec- trodes
Isotopes reported	Te ¹²⁶ Te ¹²⁸ Te ¹³⁰	Te ¹²⁵	Te ¹²² Te ¹²³ Te ¹²⁴ Te ¹²⁷	Te ¹¹⁸ Te ¹²⁰ Te ¹²⁴ Te ¹³⁰	No Te ¹¹⁸ nor Te ¹²⁷ Te ¹²⁰
Mass number	Abundances (per cent)				
120			-----		
122			2.9		
123			1.6		
124			4.5		
125		6.6	6.0		
126	Te ¹²⁸ and Te ¹³⁰ equal	20.9	19.0		
128	intensity and double that of Te ¹²⁵	36.1	32.8		
130		36.4	33.1		

	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000	1001	1002	1003	1004	1005	1006	1007	1008	1009	1010	1011	1012	1013	1014	1015	1016	1017	1018	1019	1020	1021	1022	1023	1024	1025	1026	1027	1028	1029	1030	1031	1032	1033	1034	1035	1036	1037	1038	1039	1040	1041	1042	1043	1044	1045	1046	1047	1048	1049	1050	1051	1052	1053	1054	1055	1056	1057	1058	1059	1060	1061	1062	1063	1064	1065	1066	1067	1068	1069	1070	1071	1072	1073	1074	1075	1076	1077	1078	1079	1080	1081	1082	1083	1084	1085	1086	1087	1088	1089	1090	1091	1092	1093	1094	1095	1096	1097	1098	1099	1100	1101	1102	1103	1104	1105	1106	1107	1108	1109	1110	1111	1112	1113	1114	1115	1116	1117	1118	1119	1120	1121	1122	1123	1124	1125	1126	1127	1128	1129	1130	1131	1132	1133	1134	1135	1136	1137	1138	1139	1140	1141	1142	1143	1144	1145	1146	1147	1148	1149	1150	1151	1152	1153	1154	1155	1156	1157	1158	1159	1160	1161	1162	1163	1164	1165	1166	1167	1168	1169	1170	1171	1172	1173	1174	1175	1176	1177	1178	1179	1180	1181	1182	1183	1184	1185	1186	1187	1188	1189	1190	1191	1192	1193	1194	1195	1196	1197	1198	1199	1200	1201	1202	1203	1204	1205	1206	1207	1208	1209	1210	1211	1212	1213	1214	1215	1216	1217	1218	1219	1220	1221	1222	1223	1224	1225	1226	1227	1228	1229	1230	1231	1232	1233	1234	1235	1236	1237	1238	1239	1240	1241	1242	1243	1244	1245	1246	1247	1248	1249	1250	1251	1252	1253	1254	1255	1256	1257	1258	1259	1260	1261	1262	1263	1264	1265	1266	1267	1268	1269	1270	1271	1272	1273	1274	1275	1276	1277	1278	1279	1280	1281	1282	1283	1284	1285	1286	1287	1288	1289	1290	1291	1292	1293	1294	1295	1296	1297	1298	1299	1300	1301	1302	1303	1304	1305	1306	1307	1308	1309	1310	1311	1312	1313	1314	1315	1316	1317	1318	1319	1320	1321	1322	1323	1324	1325	1326	1327	1328	1329	1330	1331	1332	1333	1334	1335	1336	1337	1338	1339	1340	1341	1342	1343	1344	1345	1346	1347	1348	1349	1350	1351	1352	1353	1354	1355	1356	1357	1358	1359	1360	1361	1362	1363	1364	1365	1366	1367	1368	1369	1370	1371	1372	1373	1374	1375	1376	1377	1378	1379	1380	1381	1382	1383	1384	1385	1386	1387	1388	1389	1390	1391	1392	1393	1394	1395	1396	1397	1398	1399	1400	1401	1402	1403	1404	1405	1406	1407	1408	1409	1410	1411	1412	1413	1414	1415	1416	1417	1418	1419	1420	1421	1422	1423	1424	1425	1426	1427	1428	1429	1430	1431	1432	1433	1434	1435	1436	1437	1438	1439	1440	1441	1442	1443	1444	1445	1446	1447	1448	1449	1450	1451	1452	1453	1454	1455	1456	1457	1458	1459	1460	1461	1462	1463	1464	1465	1466	1467	1468	1469	1470	1471	1472	1473	1474	1475	1476	1477	1478	1479	1480	1481	1482	1483	1484	1485	1486	1487	1488	1489	1490	1491	1492	1493	1494	1495	1496</
--	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	--------

Table II cont'd

Worker Year Reference	Migeotto 1936 (77)	Williams and Yuster 1946 (132)	White and Cameron 1948 (131)	Hollander et al. 1953 (54)
Instrument	Bainbridge mass spectro- scope, photo- metric detec- tion	Nier 60° mass spec- trometer, magnetic scanning, electron- ic detection	60° mass spectro- meter, magnetic scanning, electronic detector	
Ion source	Te ions excited by Hg 4359 line	Ionization of TeF ₆ gas with 100 volt electrons	Bombarding TeF ₆ gas with 30-75 volt electrons	
Isotopes reported	Te ¹²⁰ Te ¹²² Te ¹²³ Te ¹²⁴ Te ¹²⁵ Te ¹²⁶ Te ¹²⁸ Te ¹³⁰	same	same	same
Mass number	Abundances (%)			
	* Average	TeF ₃	TeF ₅	
120	1.5	0.26	0.091	0.089
122	2.9	7.05	2.49	2.46
123	1.6	2.47	0.89	0.87
124	4.5	13.3	4.63	4.61
125	6.0	20.15	7.01	6.99
126	19.0	54.2	18.72	18.71
128	32.8	92.3	31.72	31.79
130	33.1	100	34.49	34.49

*Relative abundances

in a valence state as high as +4. This results because of the different oxidation potentials for the elements. Further, similar chemical reactions of these elements have different temperature dependences, and because of their different sizes, the three elements behave differently in physical processes such as diffusion.

If all three elements can be simultaneously examined isotopically, it may be possible to separate the various physical, chemical, and biological processes which have occurred in a deposit.

THEORY

An isotope effect, or isotope fractionation, occurs when the relative abundances of the isotopes of an element in some system are altered by physical or chemical means, and is a result of slight differences in the properties of these isotopes. Chemically an effect may occur when two or more molecular species containing the element in question co-exist in chemical equilibrium (equilibrium isotope effect) or when one or more molecular species change their identity in the course of a chemical reaction (kinetic isotope effect).

The principles of isotopic fractionation are now well-known and it is possible to calculate, from a knowledge of the fundamental vibrational frequencies of the compounds, the extent to which the isotopes might be expected to fractionate in a naturally occurring process.

Looking at the members of the group VIA elements, there is a mass difference of two or a percentage mass difference of 13% in the case of the two oxygen isotopes O^{16} and O^{18} ; a mass difference of two or a percentage mass difference of 7% in the case of S^{32} and S^{34} ; a mass difference of six or percentage mass difference of 8% in the case of Se^{76} and Se^{82} ; and a mass difference of eight or percentage mass difference of 7% in the case of Te^{130} and Te^{122} .

Significant isotope effects have been found for the two oxygen, sulphur, and selenium isotopes respectively, and therefore, some fractionation of the tellurium isotopes is expected.


A. Theory of Equilibrium Isotope Effects

Although chemical equilibrium may be considered as that condition of a chemical system where the rate at which a forward reaction is taking place is exactly balanced by the rate of the reverse reaction, the theory of equilibrium isotope effects is best considered in terms of the statistical mechanics of a system of particles in dynamic equilibrium.

Urey and Rittenberg (114), (115), were the first to show by theoretical calculations that marked differences in the equilibrium constants of isotopic exchange reactions should exist. Urey and Grieff (116) applied statistical mechanics to the calculation of equilibrium exchange constants in terms of distribution functions and "summation over states" terms and this was simplified by Urey (118) and Bigeleisen and Mayer (10). Their simplification made it possible to calculate the equilibrium constants of isotopic reactions solely with a knowledge of the vibrational frequencies of the isotopic molecules. It is to be noted,

however, that such a treatment does not consider the time taken to attain equilibrium or whether there in fact exists a mechanism by which equilibrium may be attained.

An isotopic exchange reaction may be written



where the subscripts 1 and 2 refer to the light and heavy isotopes respectively and A and B are molecules having the element under consideration as a common constituent. From thermodynamics (72), the standard free energy change for the above reaction is given in terms of the equilibrium constant by

$$-RT \ln K = \Delta F^0 \quad (2)$$

or

$$-RT \ln K = aF_{A_2} + bF_{B_1} - aF_{A_1} - bF_{B_2} \quad (3)$$

In terms of the partition function "Q", the free energy can be written as (F = Gibbs free energy)

$$F = E_0 + RT \ln N - RT \ln Q \quad (4)$$

where N is Avagadro's number, E_0 is the zero point energy, and Q is the total partition function given by

$$Q = \sum_n g_n e^{-\epsilon_n/kT} \quad (5)$$

where g_n is the statistical weight factor and the summation extends over all the quantum states of energy ϵ_n . A d-fold degenerate level would count as d states.

If only one molecule is considered in unit volume instead of one mole, N is replaced by unity and the second term of (4) disappears. Substitution of (4) into (3) and simplification gives

$$K = \left\{ \left[\frac{Q_{A_2}}{Q_{A_1}} \right]^a / \left[\frac{Q_{B_2}}{Q_{B_1}} \right]^b \right\} \exp \left[\frac{aE_{oA_2} + bE_{oB_1} - aE_{oA_1} - bE_{oB_2}}{kT} \right] \quad (6)$$

Since they are determined by the electrical forces, the potential energy curves have been found to be essentially identical for isotopic molecules. Referring all energies to the minimum of the potential energy curves rather than the zero point energy simplifies (6) to

$$K = \left[\frac{Q_{A_2}}{Q_{A_1}} \right]^a / \left[\frac{Q_{B_2}}{Q_{B_1}} \right]^b \quad (7)$$

and the equilibrium constant becomes the ratios of the partition functions for the two isotopically substituted molecules.

B. Evaluation of Partition Function Ratios

The partition function of a molecule depends upon its mass, moments of inertia, fundamental vibrational frequencies, temperature, and nuclear spins of the constituent atoms. It is, however, possible to evaluate partition function ratios for isotopic molecules using only vibrational frequencies.

Following Herzberg (49) considering a volume V ,

$$Q = \frac{(2\pi MkT)^{3/2}}{h^3} \cdot V \cdot \frac{kT}{hcB} \prod_i^{3n-5} \frac{e^{-u_i/2}}{1-e^{-u_i}} \frac{1}{\sigma} \prod (2I+1) \prod \frac{(2I+1)^2}{2} \quad (8)$$

for diatomic (or linear polyatomic) molecules, and

$$Q = \frac{(2\pi MkT)^{3/2}}{h^3} \cdot V \cdot \left[\frac{\pi}{ABC} \left(\frac{kT}{hc} \right)^3 \right]^{1/2} \prod_i^{3n-6} \frac{e^{-u_i/2}}{1-e^{-u_i}} \left[\frac{1}{\sigma} \prod (2I+1) \prod \frac{(2I+1)^2}{2} \right] \quad (9)$$

for polyatomic molecules, where

n = number of atoms in the molecule

M = total molecular mass

k = Boltzman constant

A, B, C = rotational constants of the molecules, related to its principal moments of inertia, i.e.,

$$A = \frac{h}{8\pi^2 c I_A}$$

$u_i = \frac{h\nu_i}{kT}$ where ν_i is the i^{th} fundamental frequency of the molecule.

The product $\prod_i$ extends over all i fundamental frequencies of the polyatomic molecule and the bracketed terms are symmetry and nuclear spin factors. The nuclear spin factors cancel out in later expressions and have no effect on the estimation of isotope effects. The symmetry factors are discussed later.

Thus, for two isotopic polyatomic molecules,

$$\frac{Q_2}{Q_1} = \left[\frac{M_2}{M_1} \right]^{3/2} \cdot \left[\frac{I_{A_2} I_{B_2} I_{C_2}}{I_{A_1} I_{B_1} I_{C_1}} \right]^{1/2} \prod_i \frac{e^{-u_{2i}/2}}{1 - e^{-u_{2i}}} \cdot \frac{1 - e^{-u_{1i}}}{e^{-u_{1i}/2}} \quad (10)$$

Urey (118) simplified the two expressions (8) and (9) and defined new partition functions which were the equilibrium constants for exchange reactions between the compound considered and the separated atoms. Both sides of equations (8) and (9) are multiplied by $(M_1/M_2)^{3/2n}$ where M_1 and M_2 are the atomic weights of the isotopic atoms being considered and n is the number of the isotopic atoms exchanged. Further, the right sides of (8) and (9) are multiplied and divided by (u_1/u_2) and $\prod_i (u_{1i}/u_{2i})$ respectively. According to the Teller-Redlich theorem (70,87),

$$\frac{I_2}{I_1} \left[\frac{M_2}{M_1} \right]^{3/2} \left[\frac{M_1}{M_2} \right]^{3/2n} \left[\frac{u_1}{u_2} \right] = \left[\frac{A_2 B_2 C_2}{A_1 B_1 C_1} \right]^{1/2} \left[\frac{M_2}{M_1} \right]^{3/2} \left[\frac{M_1}{M_2} \right]^{3/2n} \left[\frac{u_1}{u_2} \right] \quad (11)$$

Carrying out the steps above and using (11), equation (10) becomes

$$\frac{Q_2}{Q_1} = \left[\frac{M_{2i}}{M_{1i}} \right]^{3/2} \frac{\sigma_1}{\sigma_2} \cdot \frac{u_2}{u_1} \cdot \frac{e^{-u_2/2} (1 - e^{-u_1})}{e^{-u_1/2} (1 - e^{-u_2})} \quad (12)$$

and

$$\frac{Q_2}{Q_1} = \frac{3n-6}{i} \left[\frac{M_{2i}}{M_{1i}} \right]^{3/2} \frac{\sigma_1}{\sigma_2} \frac{3n-6}{i} \frac{u_{2i}}{u_{1i}} \cdot \frac{e^{-u_2/2} (1 - e^{-u_1})}{e^{-u_1/2} (1 - e^{-u_2})} \quad (13)$$

for diatomic molecules and polyatomic molecules respectively.

The terms $(M_{2i}/M_{1i})^{3/2}$ will not affect the expression for an equilibrium constant since they will appear in both numerator and denominator. Thus the equilibrium constant can be given by

$$K = \left[\frac{Q'_{2A}}{Q'_{1A}} \right]^a / \left[\frac{Q'_{2B}}{Q'_{1B}} \right]^b \quad (14)$$

where $\frac{Q'_2}{Q'_1} = \left[\frac{M_1}{M_2} \right]^{3/2n}$. $\frac{Q_2}{Q_1}$ is the relation between the new partition function ratio and the old ratio. K can now

be calculated with only a knowledge of the vibrational frequencies of the isotopic molecules. These frequencies are obtained either by direct observation or by calculation from observations on a more abundant isotope by means of force equations.

This expression excludes the anharmonic terms in the vibrational energy, and assumes that the rotational partition functions have reached classical values. However, these have been investigated by Urey (118) and Herzberg (49) and the approximation is not greatly in error when considering the ratios of partition functions for isotopic molecules, provided the observed fundamental frequencies are used rather than the zero order frequencies and that the temperature is low enough so that only low lying vibrational states have appreciable populations. The ratio of symmetry numbers will be unity if the molecule under consideration contains only one atom of the element for which an exchange is considered or if the molecule contains more than one such atom but these atoms occupy indistinguishable positions in the molecule and are all exchanged in the reaction.

Thus, we have for diatomic molecules

$$\frac{Q_2'}{Q_1'} = \frac{u_2}{u_1} \cdot \frac{e^{-u_2/2} \cdot (1 - e^{-u_1})}{e^{-u_1/2} \cdot (1 - e^{-u_2})} \quad (15)$$

and for polyatomic molecules,

$$\frac{Q_2'}{Q_1'} = \prod_i^{\frac{3n-6}{2}} \frac{u_{2i}}{u_{1i}} \cdot \frac{e^{-u_{2i}} \cdot (1 - e^{-u_{1i}})}{e^{-u_{1i}} \cdot (1 - e^{-u_{2i}})} \quad (16)$$

These expressions have been simplified by Urey (118) and Bigeleisen and Mayer (10) for manual calculations. In the work of this thesis IBM 7040 programs were set up to calculate the exact expressions directly.

C. Kinetic Isotope Effects

The theory of isotope fractionation in kinetic chemical reactions is considered in terms of the theory of absolute reaction rates. This theory was developed by Eyring (36) and Evans and Polanyi (35) from a statistical treatment of reaction rates. Bigeleisen (9) then used their theory and collision theory to develop formulae which predict the kinetic isotope effect.

This theory is developed around two postulates. Firstly, there exists an "activated complex" in chemical equilibrium with the initial state of the reactants and intermediate between this initial state and a final state of the products. Secondly, this "activated complex" decomposes at a definite rate to form the products of the reaction.

From these postulates, the rate constant k can be written (40a) as

$$k = \frac{k'T}{h} \cdot k_T t K$$

where

k' = Boltzman constant

T = absolute temperature

h = Planck's constant

k_T = transmission coefficient

t = quantum mechanical tunnelling correction

K = equilibrium constant for the equilibrium between the reactant and the activated complex

Thus, for isotopic molecules, the ratio of rate constants becomes,

$$\frac{k_1}{k_2} = \frac{k_{T1}}{k_{T2}} \cdot \frac{t_1}{t_2} \cdot \frac{K_1}{K_2}$$

Hirschfelder and Wigner (52) have conducted a theoretical study to show that the transmission coefficients ratio is very nearly unity for systems above room temperature having a distribution in the velocities of the reacting isotopic molecules. Also, t_1 is set equal to t_2 , the assumption of which is discussed by Bigeleisen (13).

Thus we can write

$$\frac{k_1}{k_2} = \frac{K_1}{K_2}$$

From the previous discussion of equilibrium constants,

$$\frac{K_1}{K_2} = \frac{Q_2'}{Q_1'} \bigg/ \frac{Q_2^\ddagger}{Q_1^\ddagger} \quad (17)$$

where Q_1 and Q_2 refer to partition functions for the reactants, and $Q_1^\ddagger$ and $Q_2^\ddagger$ refer to partition functions for the activated complexes. The partition function ratios can be set up in the same manner as was done in part A of the theory. For the activated complex, however, we have to consider its nature. According to the theory, one of the vibrational degrees of freedom of the activated complex is non-genuine, corresponding to the cleavage (or formation) of a bond. Thus, supposing that the L^{th} vibrational mode is non-genuine, so that U_L has a pure imaginary value, the vibrational contribution to the partition function of the activated complex will then take the form

$$\frac{3n^\ddagger - 7}{i} \frac{e^{-U_i/2}}{1 - e^{-U_i}}$$

so that

$$\frac{Q_2^\ddagger}{Q_1^\ddagger} = \frac{3n^\ddagger - 7}{j} \left[\frac{M_{2j}^\ddagger}{M_{1j}^\ddagger} \right]^{3/2} \frac{U_{2L}^\ddagger}{U_{1L}^\ddagger} \prod_{i \neq L} \frac{U_{2i}^\ddagger}{U_{1i}^\ddagger} \cdot \frac{e^{-U_{2i}^\ddagger/2} (1 - e^{-U_{1i}^\ddagger})}{e^{-U_{1i}^\ddagger/2} (1 - e^{-U_{2i}^\ddagger})}$$

for polyatomic molecules.

Again, the mass ratios will cancel so we can write

$$\frac{Q_2^{\ddagger''}}{Q_1^{\ddagger''}} = \frac{U_{2L}^\ddagger}{U_{1L}^\ddagger} \prod_{i \neq L} \frac{U_{2i}^\ddagger}{U_{1i}^\ddagger} \cdot \frac{e^{-U_{2i}^\ddagger} (1 - e^{-U_{1i}^\ddagger})}{e^{-U_{1i}^\ddagger/2} (1 - e^{-U_{2i}^\ddagger})}$$

now

$$U_L = \frac{hc}{kT} \cdot \omega_L \quad \omega = \text{frequency}$$

so that

$$\frac{U_{2L}^\ddagger}{U_{1L}^\ddagger} = \frac{\omega_{2L}^\ddagger}{\omega_{1L}^\ddagger}$$

Furthermore, this "imaginary" frequency is the stretching frequency of the bond being broken (or formed). Thus, this frequency can be expressed as the square root of the inverse of the effective masses along the co-ordinate of decomposition of the activated complex. Our expression then becomes

$$\frac{Q_2^\ddagger}{Q_1^\ddagger} = \left[\frac{M_2^\ddagger}{M_1^\ddagger} \right]^{1/2} \frac{3n^\ddagger - 7}{i \neq L} \frac{U_{2i}^\ddagger}{U_{1i}^\ddagger} \cdot \frac{e^{-U_{2i}^\ddagger/2} (1 - e^{-U_{1i}^\ddagger})}{e^{-U_{1i}^\ddagger/2} (1 - e^{-U_{2i}^\ddagger})}$$

writing this as

$$\frac{Q_2^\ddagger}{Q_1^\ddagger} = \left[\frac{M_2^\ddagger}{M_1^\ddagger} \right]^{1/2} \frac{Q_2^\ddagger}{Q_1^\ddagger}$$

it can be seen that equation (17) can be written as

$$\frac{k_1}{k_2} = \left[\frac{Q_2^\ddagger}{Q_1^\ddagger} \right] \cdot \left[\frac{M_2^\ddagger}{M_1^\ddagger} \right]^{1/2} \quad (18)$$

From equation (18) it can be seen that the light molecule usually has a greater rate constant since the factor $(M_2^\ddagger/M_1^\ddagger)^{1/2}$ is always greater than unity and the reacting molecule is usually more tightly bound than the activated complex. That is

$$\frac{Q_2^\ddagger}{Q_1^\ddagger} > \frac{Q_2^\ddagger}{Q_1^\ddagger}$$

第一編 總論

第一章 總論

第一節 總論

第一節 總論

第二節 總論

第三節 總論

第四節 總論

第五節 總論

第六節 總論

第七節 總論

第八節 總論

Regarding the ratio ($M_2^\ddagger/M_1^\ddagger$), two different approaches can be made. Wolfsberg (136) gives the following reasoning. Consider a molecule, X-A-B-Y in which the A-B bond is broken. If the A-B bond is strong compared to the X-A, B-Y bonds, then the latter will shorten as the former lengthens and the reduced mass will be closer to ($\frac{AB}{A+B}$). If, however, the A-B bond is the weaker, then the reduced mass will be closer to ($M_{XA} \cdot M_{XB} / (M_{XA} + M_{XB})$) since the mass centres M_{XA} and M_{XB} are being separated.

Since very little is known about the nature of the activated complex, the rate constant cannot be evaluated specifically. However, three different approaches are feasible which follow the formalism of the theory and incorporate the idea that in the transition state, the vibrational frequencies of the activated complex are the same as those of the reactant molecule except for that corresponding to bond cleavage or formation. Approaches I and III give extreme lower and upper limits respectively that might be expected for the ratio of rate constants and approach II gives an intermediate and best description of the physical situation.

1890

1890

1890

I. When the partition function ratio term is very close to unity which occurs when the temperature is high. In the high temperature limit,

$$\frac{k_1}{k_2} = \left[\frac{M_2^\ddagger}{M_1^\ddagger} \right]^{1/2}$$

The mass ratio can either be the atomic mass ratio or mass fragment ratio.

II. Where the activated complex is identical to the starting material, with the exception that the frequency associated with the breaking bond is missing. In this case,

$$\frac{Q_2'}{Q_1'} \bigg/ \frac{Q_2^\ddagger}{Q_1^\ddagger} = \frac{Q(\omega_{2L})}{Q(\omega_{1L})}$$

where this ratio is the same as the partition function ratio for diatomic isotopic molecules with frequencies ω_{2L} and ω_{1L} , ω_L being that frequency in the starting material which is missing in the activated complex. The mass ratio is evaluated as before.

III. Where the activated complex has some structure which is strictly speaking unknown, but for which the value of the partition function ratio lies between that for the starting material (with the omission of a


frequency term) and unity. The extreme case of unity may be approached if in the transition state there is a complexing of the reacting molecule with other material, so that a complicated structure is formed in which not many of the vibrational frequencies are sensitive to isotopic substitution. In this case,

$$\frac{k_1}{k_2} = \frac{Q_2}{Q_1} \cdot \left[\frac{M_2^\ddagger}{M_1^\ddagger} \right]^{1/2}$$

These three approaches give for a particular model of the activated complex an intermediate (II) as well as extreme (I, and III) estimates for the ratio of rate constants.

D. Isotope fractionation during first order competing reactions

In general, we have the competitive isotopic reactions of the type



where subscripts 1 and 2 refer to the light and heavy isotopic species respectively. The rate constant k

...the ... of ...
...the ... of ...
...the ... of ...
...the ... of ...
...the ... of ...



...the ... of ...
...the ... of ...
...the ... of ...
...the ... of ...
...the ... of ...

...the ... of ...
...the ... of ...

...the ... of ...
...the ... of ...

...the ... of ...
...the ... of ...
...the ... of ...
...the ... of ...
...the ... of ...

...the ... of ...
...the ... of ...
...the ... of ...
...the ... of ...
...the ... of ...

indicates how fast A is converted to X.

The rate of the reaction is given in terms of the rate constant as

$$\text{Rate} = kC_A C_B$$

where k = the rate constant

C_A = concentration of reactant A

C_B = concentration of reactant B

A first order reaction is one where the time dependence of product formation can be written as

$$\frac{dX}{dt} = k(A_0 - X)B$$

where A_0 is the initial concentration of the reactant.

Thus, for the competing reactions mentioned,

$$\frac{dX_1}{dt} = k_1(A_{01} - X_1)B$$

$$\frac{dX_2}{dt} = k_2(A_{02} - X_2)B$$

At $t = 0$, $X_1 = X_2 = 0$, so upon integrating,

$$\frac{k_1}{k_2} = \frac{\log \left[\frac{A_{01}}{A_{01} - X_1} \right]}{\log \left[\frac{A_{02}}{A_{02} - X_2} \right]}$$

If we designate the fraction of molecules which have reacted as

$$f = \frac{X_1 + X_2}{A_{01} + A_{02}} = \frac{X_1(1 + X_2/X_1)}{A_{01}(1 + A_{02}/A_{01})}$$

and the integrated isotopic composition of the product as

$$r = \frac{X_2/X_1}{A_{02}/A_{01}}$$

we can write

$$\frac{k_1}{k_2} = \frac{\log \left[1 - f \frac{1 + A_{02}/A_{01}}{1 + X_2/X_1} \right]}{\log \left[1 - rf \frac{1 + A_{02}/A_{01}}{1 + X_2/X_1} \right]} \quad (19)$$

Thus, the ratio of the rate constants for competitive isotopic reactions is obtained from the determination of isotopic ratios in the initial reactant A_{02}/A_{01} and in the product X_2/X_1 after fraction f of the molecules have reacted.

THE UNIVERSITY OF CHICAGO PRESS

54 EAST LAKE STREET, CHICAGO, ILL. 60601

THE UNIVERSITY OF CHICAGO PRESS
54 EAST LAKE STREET, CHICAGO, ILL. 60601

THE UNIVERSITY OF CHICAGO PRESS
54 EAST LAKE STREET, CHICAGO, ILL. 60601

THE UNIVERSITY OF CHICAGO PRESS
54 EAST LAKE STREET, CHICAGO, ILL. 60601

THE UNIVERSITY OF CHICAGO PRESS
54 EAST LAKE STREET, CHICAGO, ILL. 60601

THE UNIVERSITY OF CHICAGO PRESS
54 EAST LAKE STREET, CHICAGO, ILL. 60601

THE UNIVERSITY OF CHICAGO PRESS
54 EAST LAKE STREET, CHICAGO, ILL. 60601

THE UNIVERSITY OF CHICAGO PRESS
54 EAST LAKE STREET, CHICAGO, ILL. 60601

THE UNIVERSITY OF CHICAGO PRESS
54 EAST LAKE STREET, CHICAGO, ILL. 60601

THE UNIVERSITY OF CHICAGO PRESS
54 EAST LAKE STREET, CHICAGO, ILL. 60601

THE UNIVERSITY OF CHICAGO PRESS
54 EAST LAKE STREET, CHICAGO, ILL. 60601

THEORETICAL CALCULATIONS

A. Calculation of equilibrium exchange constants

As was noted, the extent to which one would expect isotopic fractionation to take place in exchange reactions is found from the partition function ratios. Thus, the partition function ratios have been calculated for various molecules containing Te^{130} and Te^{122} .

Appendix I contains calculations of vibrational frequencies for some Te^{130} and Te^{122} -containing compounds for which molecular data such as symmetry type and fundamental vibrational frequencies were available. For the diatomic species, anharmonicity corrections were made. Since Te^{130} is the most abundant isotope, and also since use of this mass number gave results closest to the observed, it was assumed that fundamental frequencies applied to compounds containing this isotope. The frequencies of the Te^{122} -containing molecules were then calculated by means of force field equations from these observed values.

In most cases, sets of equations describing the force fields were accompanied by force constants already calculated. However, in the case of H_2Te and TeO_3^- the force constants had not been previously calculated so

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

these were calculated by substituting the known fundamental frequencies into a set of force equations describing the molecule.

From the isotopic vibrational frequencies, the partition function ratios at various temperatures were calculated using equations (15) and (16) for diatomic and polyatomic molecules respectively. The equilibrium exchange constants for possible isotopic exchange reactions were then calculated using equation (14). Tables III and IV give the partition function ratios and equilibrium exchange constants respectively. Looking at Table IV, the values in the body of the table are the equilibrium constants for exchange reactions between the compounds listed along the top and the compounds listed down the left-hand column calculated for the temperatures listed in the far right-hand column. A constant greater than unity means that the heavier isotope is concentrated in the compound in the left-hand column.

For example, consider TeF_6 in the left-hand column, and TeO at the top of the table. The value 1.0234 is found at 20°C in the table where the TeO column and TeF_6 row intersect. This is the equilibrium constant for the reaction:

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

...and the ...

Table III

 Q_2'/Q_1' for Te-containing molecules

$^{\circ}\text{C}$	GeTe	PbTe	SnTe	TeO	SiTe	TeTe	HTe	TeF ₆	TeBr ₄	SCTe	TeO ₃ [−]	H ₂ Te
0	1.00270	1.00205	1.00235	1.00409	1.00278	1.00230	1.00114	1.03067	1.00467	1.00833	1.01398	1.00241
10	1.00252	1.00191	1.00219	1.00386	1.00261	1.00215	1.00110	1.02882	1.00435	1.00779	1.01313	1.00230
20	1.00236	1.00178	1.00205	1.00364	1.00244	1.00201	1.00105	1.02713	1.00406	1.00731	1.01236	1.00220
30	1.00221	1.00167	1.00192	1.00344	1.00230	1.00188	1.00101	1.02559	1.00380	1.00686	1.01165	1.00211
40	1.00208	1.00157	1.00181	1.00326	1.00216	1.00177	1.00097	1.02416	1.00357	1.00646	1.01100	1.00202
50	1.00196	1.00147	1.00170	1.00309	1.00204	1.00166	1.00093	1.02285	1.00335	1.00609	1.01040	1.00193
60	1.00184	1.00139	1.00160	1.00293	1.00193	1.00156	1.00090	1.02164	1.00316	1.00575	1.00985	1.00186
70	1.00174	1.00131	1.00151	1.00279	1.00182	1.00148	1.00086	1.02052	1.00298	1.00544	1.00933	1.00178
80	1.00165	1.00124	1.00143	1.00265	1.00173	1.00140	1.00083	1.01949	1.00281	1.00515	1.00886	1.00172
90	1.00156	1.00117	1.00135	1.00252	1.00164	1.00132	1.00080	1.01853	1.00266	1.00489	1.00842	1.00165
100	1.00148	1.00111	1.00128	1.00241	1.00155	1.00125	1.00077	1.01764	1.00252	1.00464	1.00802	1.00159
200	1.00093	1.00069	1.00080	1.00157	1.00099	1.00078	1.00056	1.01136	1.00157	1.00295	1.00515	1.00113
300	1.00064	1.00047	1.00055	1.00110	1.00068	1.00053	1.00042	1.00790	1.00107	1.00204	1.00358	1.00085
400	1.00046	1.00034	1.00040	1.00081	1.00050	1.00039	1.00033	1.00579	1.00078	1.00149	1.00262	1.00065
500	1.00035	1.00026	1.00030	1.00062	1.00038	1.00029	1.00026	1.00443	1.00059	1.00113	1.00200	1.00052
600	1.00028	1.00020	1.00024	1.00049	1.00030	1.00023	1.00022	1.00349	1.00046	1.00089	1.00158	1.00042
700	1.00022	1.00016	1.00019	1.00040	1.00024	1.00019	1.00018	1.00282	1.00037	1.00072	1.00128	1.00035
800	1.00018	1.00014	1.00016	1.00033	1.00020	1.00015	1.00015	1.00233	1.00031	1.00059	1.00105	1.00029
900	1.00015	1.00011	1.00013	1.00027	1.00017	1.00013	1.00013	1.00195	1.00026	1.00050	1.00088	1.00025
1000	1.00013	1.00010	1.00011	1.00023	1.00014	1.00011	1.00011	1.00165	1.00022	1.00042	1.00075	1.00021

Table IV. Equilibrium Constants for Te^{130} -- Te^{122} Exchange Reactions

	$\text{Te}^{130}\text{F}_6$ $\text{Te}^{122}\text{F}_6$	$\text{Te}^{130}\text{O}_3$ $\text{Te}^{122}\text{O}_3$	SCTe^{130} SCTe^{122}	$\text{Te}^{130}\text{Br}_4$ $\text{Te}^{122}\text{Br}_4$	Te^{130} Te^{122}	SiTe^{130} SiTe^{122}	GeTe^{130} GeTe^{122}	$\text{H}_2\text{Te}^{130}$ $\text{H}_2\text{Te}^{122}$	SnTe^{130} SnTe^{122}	$\text{Te}^{130}\text{Te}^{122}$ $\text{Te}^{130}\text{Te}^{122}$	PbTe^{130} PbTe^{122}	HTe^{130} HTe^{122}	$T^\circ\text{C}$
$\text{Te}^{130}\text{F}_6$ $\text{Te}^{122}\text{F}_6$	1.0000	1.0165	1.0222	1.0259	1.0265	1.0298	1.0279	1.0281	1.0283	1.0283	1.0286	1.0294	0
		1.0146	1.0197	1.0230	<u>1.0234</u>	1.0246	1.0247	1.0249	1.0250	1.0251	1.0253	1.0261	20
		1.0123	1.0167	1.0194	1.0197	1.0208	1.0208	1.0209	1.0211	1.0212	1.0213	1.0219	50
		1.0095	1.0129	1.0151	1.0152	1.0161	1.0161	1.0160	1.0163	1.0164	1.0165	1.0169	100
$\text{Te}^{130}\text{O}_3$ $\text{Te}^{122}\text{O}_3$		1.0000	1.0056	1.0093	1.0098	1.0112	1.0112	1.0115	1.0116	1.0117	1.0119	1.0128	0
			1.0050	1.0083	1.0087	1.0099	1.0100	1.0101	1.0103	1.0103	1.0106	1.0113	20
			1.0043	1.0070	1.0073	1.0083	1.0084	1.0085	1.0087	1.0087	1.0089	1.0095	50
			1.0034	1.0055	1.0056	1.0065	1.0065	1.0064	1.0067	1.0068	1.0069	1.0072	100
SCTe^{130} SCTe^{122}			1.0000	1.0036	1.0042	1.0055	1.0056	1.0059	1.0060	1.0060	1.0063	1.0072	0
				1.0032	1.0037	1.0049	1.0049	1.0051	1.0052	1.0053	1.0055	1.0063	20
				1.0027	1.0030	1.0040	1.0041	1.0042	1.0044	1.0044	1.0046	1.0052	50
				1.0021	1.0022	1.0031	1.0032	1.0030	1.0034	1.0034	1.0035	1.0039	100
$\text{Te}^{130}\text{Br}_4$ $\text{Te}^{122}\text{Br}_4$				1.0000	1.0006	1.0019	1.0020	1.0023	1.0023	1.0024	1.0026	1.0035	0
					1.0004	1.0016	1.0017	1.0019	1.0020	1.0020	1.0023	1.0030	20
					1.0003	1.0013	1.0014	1.0014	1.0016	1.0017	1.0019	1.0024	50
					1.0001	1.0010	1.0010	1.0009	1.0012	1.0013	1.0014	1.0017	100
Te^{130}O Te^{122}O					1.0000	1.0013	1.0014	1.0017	1.0017	1.0018	1.0020	1.0029	0
						1.0012	1.0013	1.0014	1.0016	1.0016	1.0019	1.0026	20
						1.0010	1.0011	1.0012	1.0014	1.0014	1.0016	1.0022	50
						1.0009	1.0009	1.0008	1.0011	1.0012	1.0013	1.0016	100
SiTe^{130} SiTe^{122}						1.0000	1.0001	1.0004	1.0004	1.0005	1.0007	1.0016	0
							1.0001	1.0002	1.0004	1.0004	1.0007	1.0014	20
							1.0001	1.0001	1.0003	1.0004	1.0006	1.0011	50
							1.0001	1.0000	1.0003	1.0003	1.0004	1.0008	100

Table IV Cont'd

	$\frac{\text{GeTe}^{130}}{\text{GeTe}^{122}}$	$\frac{\text{H}_2\text{Te}^{130}}{\text{H}_2\text{Te}^{122}}$	$\frac{\text{SnTe}^{130}}{\text{SnTe}^{122}}$	$\frac{\text{Te}^{130}\text{Te}^{130}}{\text{Te}^{130}\text{Te}^{122}}$	$\frac{\text{PbTe}^{130}}{\text{PbTe}^{122}}$	$\frac{\text{HTe}^{130}}{\text{HTe}^{122}}$	T°C
$\frac{\text{GeTe}^{130}}{\text{GeTe}^{122}}$	1.0000	1.0003 1.0002 1.0000 1.0000	1.0005 1.0003 1.0003 1.0002	1.0004 1.0003 1.0003 1.0002	1.0006 1.0006 1.0005 1.0004	1.0016 1.0013 1.0010 1.0007	0 20 50 100
$\frac{\text{H}_2\text{Te}^{130}}{\text{H}_2\text{Te}^{122}}$		1.0000	1.0001 1.0001 1.0002 1.0003	1.0004 1.0002 1.0003 1.0003	1.0004 1.0004 1.0005 1.0005	1.0013 1.0011 1.0010 1.0003	0 20 50 100
$\frac{\text{SnTe}^{130}}{\text{SnTe}^{122}}$			1.0000	1.0000 1.0000 1.0000 1.0000	1.0003 1.0003 1.0002 1.0002	1.0012 1.0010 1.0008 1.0005	0 20 50 100
$\frac{\text{Te}^{130}\text{Te}^{130}}{\text{Te}^{130}\text{Te}^{122}}$				1.0000	1.0002 1.0002 1.0002 1.0001	1.0012 1.0010 1.0007 1.0005	0 20 50 100
$\frac{\text{PbTe}^{130}}{\text{PbTe}^{122}}$					1.0000	1.0009 1.0007 1.0005 1.0003	0 20 50 100
$\frac{\text{HTe}^{130}}{\text{HTe}^{122}}$						1.0000	

卷之四

四


四

四

四

四

四



This means that provided the reaction is possible, a 2.3% tellurium isotope effect is predicted and the TeF_6 species will be enriched by 2.3% in the heavier isotope.

B. Calculation of ratios of isotopic rate constants

As mentioned in the theory, intermediate values as well as upper and lower extremes could be determined for the ratio of kinetic rate constants. The lower limit reached at high temperatures is where

$$\frac{k_1}{k_2} = \left[\frac{M_2^\ddagger}{M_1^\ddagger} \right]^{1/2}$$

and $(M_2^\ddagger/M_1^\ddagger)^{1/2}$ represents the effective mass of the complex along the reaction co-ordinate. If this is simply the reduced mass of the atoms whose bond is directly involved in the reaction, $M_2^\ddagger$ and $M_1^\ddagger$ are the isotopic reduced masses of the molecules. Table V gives the minimum value of the rate constant ratio to be expected in a kinetic isotope reaction in which the bond is broken between the diatomic molecules listed. These are temperature independent.

Minimum Value of k_{130}/k_{122}						
GeTe	PbTe	SnTe	TeO	SiTe	TeTe	HTe
1.012	1.020	1.016	1.004	1.006	1.016	1.000

Table V

Considering now the reduction of potassium tellurite, and suppose the rate determining step is the cleavage of a Te - O bond:

- I. The high temperature limit for k_1/k_2 is the mass term 1.004 if atom fragments are considered or 1.002 if mass fragments in the "di-fragmental molecule" $\text{TeO}_2 - \text{O}$ are used.
- II. The intermediate value and closest to the physical situation is $\left[\frac{Q_2'}{Q_1'} / \frac{Q_2^\ddagger}{Q_1^\ddagger} \right] \cdot \left[\frac{M_2^\ddagger}{M_1^\ddagger} \right]^{1/2}$ where $\frac{Q_2^\ddagger}{Q_1^\ddagger}$ is simply that for $\text{TeO}_3^\ddagger$ with the frequency contribution of the Te - O bond omitted. This mode is most probably one component of the doubly degenerate E type asymmetric stretching vibration ω_3 .

Table 1: Summary of the data						
Year	Q1	Q2	Q3	Q4	Q5	Q6
2018	1.2	1.5	1.8	2.1	2.4	2.7
2019	1.3	1.6	1.9	2.2	2.5	2.8
2020	1.4	1.7	2.0	2.3	2.6	2.9
2021	1.5	1.8	2.1	2.4	2.7	3.0
2022	1.6	1.9	2.2	2.5	2.8	3.1

The data shows a consistent upward trend in the values across all quarters from 2018 to 2022. The values for each quarter are as follows:

Q1: 1.2 (2018), 1.3 (2019), 1.4 (2020), 1.5 (2021), 1.6 (2022)

Q2: 1.5 (2018), 1.6 (2019), 1.7 (2020), 1.8 (2021), 1.9 (2022)

Q3: 1.8 (2018), 1.9 (2019), 2.0 (2020), 2.1 (2021), 2.2 (2022)

Q4: 2.1 (2018), 2.2 (2019), 2.3 (2020), 2.4 (2021), 2.5 (2022)

Q5: 2.4 (2018), 2.5 (2019), 2.6 (2020), 2.7 (2021), 2.8 (2022)

Q6: 2.7 (2018), 2.8 (2019), 2.9 (2020), 3.0 (2021), 3.1 (2022)

The overall trend indicates a steady increase in the values over the five-year period, with each quarter showing a similar growth pattern.

$$\frac{\frac{Q_2'}{\ddagger}}{\frac{Q_1'}{\ddagger}} \bigg/ \frac{\frac{Q_2^{\ddagger'}}{\ddagger}}{\frac{Q_1^{\ddagger'}}{\ddagger}} = \begin{array}{ll} 1.0041 & \text{at } 0^\circ\text{C} \\ 1.0037 & \text{at } 20^\circ\text{C} \\ 1.0031 & \text{at } 50^\circ\text{C} \end{array}$$

The mass ratio is as determined in part I for either atomic or mass fragments.

III. The upper limit for k_1/k_2 is given by

$$\frac{k_1}{k_2} = \frac{Q(\text{Te}^{130}\text{O}_3^=)}{Q(\text{Te}^{122}\text{O}_3^=)} \cdot \left[\frac{\frac{\ddagger}{M_2}}{\frac{\ddagger}{M_1}} \right]^{1/2}$$

The results of the three methods of calculation are summarized in Table VI. All these simple models give a regular decrease in the value of k_1/k_2 with increase in temperature.

Table VI

Summary of the Calculated Ratios of Rate Constants for the Reduction of K_2TeO_3

T°C	k_1/k_2					
	Low		Intermediate		High	
	Mass	Atom	Mass	Atom	Mass	Atom
0	1.002	1.004	1.00615	1.00765	1.0160	1.0175
10	1.002	1.004	1.00590	1.00740	1.0152	1.0167
20	1.002	1.004	1.00568	1.00718	1.0144	1.0159
30	1.002	1.004	1.00547	1.00697	1.0137	1.0152
40	1.002	1.004	1.00528	1.00678	1.0130	1.0145
50	1.002	1.004	1.00511	1.00661	1.0124	1.0139
60	1.002	1.004	1.00494	1.00644	1.0119	1.0134
70	1.002	1.004	1.00479	1.00629	1.0114	1.0129
80	1.002	1.004	1.00465	1.00615	1.0109	1.0124
90	1.002	1.004	1.00452	1.00602	1.0104	1.0119
100	1.002	1.004	1.00440	1.00590	1.0100	1.0115
200	1.002	1.004	1.00355	1.00505	1.0072	1.0087
300	1.002	1.004	1.00308	1.00458	1.0056	1.0071
400	1.002	1.004	1.00279	1.00429	1.0046	1.0061
500	1.002	1.004	1.00261	1.00410	1.0040	1.0055
600	1.002	1.004	1.00248	1.00397	1.0036	1.0051
700	1.002	1.004	1.00239	1.00388	1.0033	1.0048
800	1.002	1.004	1.00231	1.00381	1.0031	1.0045
900	1.002	1.004	1.00227	1.00376	1.0029	1.0044
1000	1.002	1.004	1.00223	1.00372	1.0027	1.0042

EXPERIMENTAL

In investigating the isotopes of tellurium, tellurium samples were examined to see if the isotope ratio could be altered by natural and laboratory processes. Tellurium was extracted from natural telluride samples taken from various geographical locations. A controlled chemical reduction of K_2TeO_3 to elemental tellurium was carried out in the laboratory and attempts made to measure the kinetic isotope effect. K_2TeO_3 was also partially reduced to elemental tellurium by micro-organisms and the isotope composition of the products and reactant compared.

The collected elemental tellurium samples were prepared for mass spectrometric analysis by converting them to tellurium hexafluoride (TeF_6). This method was chosen because of the simplicity of the TeF_6 spectrum (there is only one stable fluorine isotope), the stability of the gas, and because of its ease of preparation. H_2Te could also be used, but the gas is unstable and its mass spectrum is quite complicated by overlapping of the ion species, eg., Te^{130} and $Te^{128}H_2^+$ both at mass 130. Computations on such a spectrum are tedious and subject to considerable error.

A. Extraction of Tellurium from Natural Samples

In 1832 it was discovered by P. Berthier that tellurium could be precipitated from raw materials with sulphur dioxide and hydrochloric acid (76). However, the extraction of pure elemental tellurium was hampered by the presence of selenium which is closely related to tellurium being of the same chemical group. Keller (58) in 1897 showed that selenium and tellurium could be separated with sulphur dioxide in differing acidities of hydrochloric acid, and Gooch and Peirce in 1898 (42) were able to separate tellurium from selenium on the basis of the different volatilities of their tetra-bromides, but this was somewhat inefficient.

To further add to the difficulty, certain heavy metals are partially precipitated by sulphur dioxide (58), and combinations of bismuth, antimony, and selenium with tellurium make separations even more complicated because of mutual interferences (88). Hillebrand and co-workers in 1953 (51) gave the different hydrochloric acid concentrations for precipitation of each of selenium and tellurium with sulphurous acid. However, they dried both elements in air at 110°C ; whereas, in the same year Duval showed that tel-

lurium oxidizes at temperatures above 40°C (30). Then in 1960, a method was described by Reed (88) by which pure elemental tellurium could be separated from all other elements. This method was employed in this investigation.

About 0.5 grams of sample are powdered and dissolved in 10 ml. of nitric acid and evaporated to dryness. The residue is dissolved in 100 ml. of concentrated hydrochloric acid with 25 ml. sulphurous acid. This step precipitates any selenium present, which is then filtered out.

To the filtrate, 2 grams of tartaric acid are added to prevent any antimony from hydrolyzing. Water is then added until the hydrochloric acid concentration is 3M. To speed up the reaction, 10 ml. of hydrazine hydrochloride solution is added and the tellurium is then precipitated with 25 ml. of sulphurous acid. After boiling for 1 or 2 minutes, the tellurium is separated from the supernatant solution by centrifuging. This is then washed and re-centrifuged several times to remove soluble ions.

The method works as follows: the nitric acid acts as an oxidizing agent producing H_2TeO_3 . The sulphurous

acid then reacts according to the equation



(The selenium was precipitated in like manner in the concentrated hydrochloric acid solution.)

According to the literature, only gold and silver would interfere in the selenium-tellurium determination if present, but the gold not being dissolved in nitric acid can be filtered out, and any silver is easily separated as chloride (58).

The efficiency of this method was tested. The precipitated tellurium was filtered through a fritted crucible, washed with 0.2N hydrochloric acid and finally water. It was dried by placing the crucible and tellurium in a desiccator at 20 mm. pressure or less at room temperature over anhydrous calcium sulphate for two hours. Of 1346.5 mg. of sample, 1323.3 mg. or 98.28% was recovered.

B. Chemical reduction of K_2TeO_3

In order to study kinetic isotope effects, the desirable conditions for the experiment are:

1. a gaseous reaction or reaction in solution,

2. a method of halting the reaction at any point, and

3. the ability to recover the reduced product.

A solution or gaseous reaction is preferable since larger isotope effects are found under these conditions. Very little isotope fractionation is expected in a solid phase reaction where the process proceeds layer by layer.


The second condition is necessary if the kinetic effect is to be measured. The reaction must be stopped and the isotope ratios determined in the unreduced reactant and the reduced product. Therefore, a chemical reduction of K_2TeO_3 in solution was chosen for study.

Halting the reaction is possible in either of two ways. A method of complete reduction can be employed and the reaction quickly halted by neutralization at the required time; or, only a percentage of the reducing agent required for complete reduction could be used. Halting by neutralization has disadvantages in that much time and many trial runs are required to determine the exact time to be allowed for the desired reduction to take place.

Furthermore, in the case of tellurium, using the hydrochloric acid-sulphur dioxide method for instance, the

reaction takes several days for complete reduction unless hydrazine-hydrochloride is used, in which case the reduction takes place too rapidly to control. Therefore, 5% - 10% reductions were affected by adding only 5% - 10% of the stoichiometrically required reducing agent.

Using the reduction equation



0.65 grams of H_2SO_3 are required for the complete reduction of one gram of K_2TeO_3 . For a 10% reduction, 10% of 0.65 grams or 65 mg. are required per gram of K_2TeO_3 . Six grams of K_2TeO_3 were reduced 10% at room temperature.

The precipitated tellurium was separated from the supernatant solution by centrifuging, was washed, re-centrifuged to remove soluble ions, and then dried at 35°C .

C. Reduction of K_2TeO_3 by *Scopulariopsis brevicaulis*

That micro-organisms are capable of reducing a complex tellurium compound to elemental tellurium has been known for several years. As early as 1914, W. E. King and L. Davis observed that nearly all of the more common micro-organisms reacted with potassium tellurite

to form "black compounds" depending on reduction of the tellurite (59). T. L. Ch'in in 1938 discovered that living colonies of pathogenic fungi could be identified by the "black color" produced upon injection of potassium tellurite (18). A year later, Bird and Challenger (13) discovered that upon ingestion of potassium tellurite, Scopulariopsis brevicaulis (*Penicillium brevicaulis*) reduced the tellurite to black elemental tellurium and evolved di-methyl telluride. Other micro-organisms were later discovered to reduce potassium tellurite to elemental tellurium such as Corynebacterium diphtheria by Morton and Anderson (79) in 1941, Mycobacterium tuberculosis by L. Sula in 1947 (102), and living tissues (bacteria, yeast, human leucocytes) by M. Wachstein in 1949 (125). Others can also be found in later literature.

To examine the isotopic variations by such a reduction, it was decided that Scopulariopsis brevicaulis should be used because it is non-pathogenic. Also, a culture of the organisms was readily obtainable from Dr. Carmichael of The Provincial Laboratory of Public Health, University of Alberta. Furthermore, growth of the culture is quite simple and may be carried out at room temperature.

The first of these is the fact that the...
...the second is the fact that the...
...the third is the fact that the...
...the fourth is the fact that the...
...the fifth is the fact that the...
...the sixth is the fact that the...
...the seventh is the fact that the...
...the eighth is the fact that the...
...the ninth is the fact that the...
...the tenth is the fact that the...
...the eleventh is the fact that the...
...the twelfth is the fact that the...
...the thirteenth is the fact that the...
...the fourteenth is the fact that the...
...the fifteenth is the fact that the...
...the sixteenth is the fact that the...
...the seventeenth is the fact that the...
...the eighteenth is the fact that the...
...the nineteenth is the fact that the...
...the twentieth is the fact that the...

1. Reduction of K_2TeO_3 by culture on liquid medium

The culture was grown on a liquid medium (Trypticase Soy Broth)* in an aerating flask. 500 ml. of TSB was inoculated with S. brevicaulis. The air was sterilized by passing it through H_2SO_4 , sterile cotton wool, then a 1:1000 solution of mercuric chloride, and again through sterile cotton wool (17). After passing through the culture flask, it was aspired through an alcoholic iodine solution to form di-methyl telluride di-iodide, $(CH_3)_2TeI_2$ (16) from any di-methyl telluride $(CH_3)_2Te$ evolved from the culture.


The culture was allowed to grow until a good growth covered the medium surface (12 days). It was then inoculated with 1.5 grams of potassium tellurite in solution, and samples were taken every 12 hours until enough of the tellurite had been reduced to yield sufficient tellurium for analysis, which took 72 hours. At this point, 10 cc. samples were taken every 6 hours and the tellurium extracted from the broth as outlined later. Finally, after 100 hours, the yield of reduced tellurium was too small for analysis and the experiment was halted. The tellurium was then extracted from the yet unreduced tellurite, the broth, and from the di-methyl telluride di-iodide compound.

*This broth was prepared by mixing 30 grams Trypticase Soy Broth powder from Baltimore Biological Company with 1 litre of water. Hitherto this will be referred to as TSB.

2. Extraction of elemental tellurium reduced from K_2TeO_3 in the biological investigation

In order to obtain only the tellurium reduced by the bacteria for analysis, it had to be separated from the rest of the broth which still contained unreduced potassium tellurite in solution and which would also be reduced by the extraction procedure. Thus, the 10 cc. sample was washed with water and filtered to separate the tellurium reduced by the bacteria from the unreduced tellurite solution. (Elemental tellurium is insoluble in water.) The filter paper was then dissolved in nitric acid and the method of extraction carried out as outlined on page 72. The filtrate was treated with concentrated hydrochloric acid and sulphur dioxide and let stand for three hours. After this, the hydrochloric acid concentration was brought to 3M, 10 ml. of hydrazine hydrochloride was added, and sulphur dioxide was bubbled through which precipitated any tellurium present. Thus, at any sampling time the bacteriologically-reduced tellurium and tellurium representative of the unreduced tellurite were available for analysis.

The tellurium was separated from the di-methyl telluride di-iodide by a method outlined by Vernon (122). Tellurium and methyl iodide react according to the equation



... ..
... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

Thus, the di-methyl telluride di-iodide was distilled in a sulphuric acid bath at 100°C evolving methyl iodide leaving a residue of elemental tellurium which was recovered.

D. Preparation of TeF_6

As stated, TeF_6 gas was chosen as the form into which the tellurium samples were converted for mass spectrometric analysis. Fluorination of elemental tellurium can result in the production of both TeF_4 and TeF_6 . According to J. H. Junkins et al (56), TeF_6 is formed at a temperature of 200°C in the presence of excess F_2 ; whereas, TeF_4 is formed at a temperature of 180°C with excess Te. In the method used, fluorine was present in a large excess and this resulted in a quantitative conversion of tellurium to TeF_6 at 200°C .

The fluorine line built for this purpose is shown in figure 5. The fluorine source was a commercial cylinder of fluorine obtained from Matheson Company of New Jersey. Many precautions are required since fluorine is most reactive and will react with everything except inert gases and already-fluorinated compounds. The reaction of fluorine with some metals such as copper and monel is retarded after an initial fluoride coating is formed so these metals can be safely used for construction of a fluorine line. Therefore, the tellurium was fluorinated and the TeF_6 trapped in a monel line.

The first part of the paper is devoted to a discussion of the
general principles of the theory of the structure of the
crystal lattice and the role of the defects in the process of
deformation.

The second part of the paper is devoted to a discussion of the
experimental results obtained in the study of the structure of the
crystal lattice and the role of the defects in the process of
deformation. The results are presented in the form of a series of
graphs and tables. The first graph shows the dependence of the
dislocation density on the degree of deformation. The second graph
shows the dependence of the dislocation density on the temperature.
The third graph shows the dependence of the dislocation density on
the strain rate. The fourth graph shows the dependence of the
dislocation density on the grain size. The fifth graph shows the
dependence of the dislocation density on the type of deformation.
The sixth graph shows the dependence of the dislocation density on
the type of material. The seventh graph shows the dependence of the
dislocation density on the type of defect. The eighth graph shows the
dependence of the dislocation density on the type of impurity.

The third part of the paper is devoted to a discussion of the
theoretical results obtained in the study of the structure of the
crystal lattice and the role of the defects in the process of
deformation. The results are presented in the form of a series of
graphs and tables. The first graph shows the dependence of the
dislocation density on the degree of deformation. The second graph
shows the dependence of the dislocation density on the temperature.
The third graph shows the dependence of the dislocation density on
the strain rate. The fourth graph shows the dependence of the
dislocation density on the grain size. The fifth graph shows the
dependence of the dislocation density on the type of deformation.
The sixth graph shows the dependence of the dislocation density on
the type of material. The seventh graph shows the dependence of the
dislocation density on the type of defect. The eighth graph shows the
dependence of the dislocation density on the type of impurity.




FIG. 5

FLUORINE LINE

Application Code



Dry TeF_6 is fairly non-reactive so that after being purified several times to remove any unreacted fluorine, the TeF_6 could be collected into glass break-seal tubes. All glass stopcocks were greased with Kel-f#90 grease since violent reactions could occur with ordinary greases upon contact with fluorine.

Safety precautions included locating both the fluorine cylinder and the entire fluorine line under a fume hood, and the wearing of a rubber coat, polyethylene gloves, and face mask during fluorinations.

The fluorination procedure is as follows. The tellurium sample is placed in a monel boat and placed in the furnace which is heated to 200°C . With V_4 closed, V_5 , V_6 , and V_7 open, the line is evacuated back as far as V_3 . After pumping on the sample to dry it thoroughly and testing the line for vacuum tightness, T_1 is cooled with liquid oxygen, valves V_5 and V_6 are closed and V_2 , V_3 , and V_4 are opened to admit a small flow of nitrogen through the line controlled by use of a flowmeter. Trap T_2 is now cooled with liquid oxygen.

With the needle valve of the fluorine cylinder closed, the main valve is opened and closed, trapping a small volume of fluorine between these valves. V_1 is then opened and the needle valve opened allowing the fluorine to pass through the line. The fluorine-nitrogen mixture is passed through a NaF trap to remove HF. T_1 freezes out any remaining HF and

water vapour present, and the TeF_6 formed is trapped in T_2 . The flow of nitrogen is maintained for a further 30 minutes to ensure scavenging of elemental fluorine from the system.

After scavenging, V_3 and V_4 are closed and the liquid oxygen dewar removed from T_2 . This vapourizes the frozen TeF_6 and releases any F_2 which might have been trapped by the TeF_6 molecules. The TeF_6 is again frozen down in T_2 and V_6 is opened to evacuate the line. This is repeated several times to ensure that no F_2 is present before admitting the TeF_6 to the glass line.

With V_6 and V_7 closed, V_5 is opened and the liquid oxygen dewar moved from T_2 to the break-seal tube. After the sample is frozen down, V_5 is closed and the liquid oxygen is replaced by a liquid-solid alcohol bath. V_7 is then opened allowing a final evacuation of the break-seal tube to remove any traces of silicon fluorides formed. V_7 is then closed and the break-seal tube removed.


E. Mass Spectrometric Analysis

In order to determine the $\text{Te}^{130}/\text{Te}^{122}$ ratio, the TeF_6 gas was analyzed in a mass spectrometer using a 12-inch radius 90° magnetic analyzer to separate ion currents of different masses. The TeF_6 molecules were ionized by an electron beam travelling normally to the inlet flow direction. The ions formed were then linearly accelerated by a 3500-volt

potential difference and subsequently separated in the magnetic field. The ion currents are introduced in turn into the collector slit by magnetic scanning. Each ion current is amplified and displayed as a peak on a continuous recording chart (see Fig. 6). Since fluorine has only one stable isotope, F^{19} , the spectra yield directly the abundances of the tellurium isotopes.

In single collection with the tracing of peaks, a realizable precision is 0.1 percent. This limitation results because the peaks are traced at different times and so fluctuations in the mass spectrometer source limit the precision. The precision can be improved by the simultaneous collection of two ion beams containing the isotopes of interest and making a direct ratio measurement of the two ion currents. Fluctuations in the ion production effect both ion currents in the same way and so the ratio remains unaltered. By introducing a standard and an unknown sample alternately into the mass spectrometer by a magnetic valve system (McKinney et al, Wanless and Thode (75)) a precision of 0.01 percent can be obtained in comparing isotope ratios of different samples.

In the ionization process, TeF_6 was split into many singly and doubly charged ion species as listed in Table VII. The TeF_5^+ ion species are the most abundant and so ion currents of masses 217 ($Te^{122}F_5^+$) and 225 ($Te^{130}F_5^+$) were simul-



taneously collected with two slits in the collector. A narrow slit (0.024") was used to collect the mass 217 ion current while a wide slit (0.040") was used for the mass 225 ion current. With the former, the resolution of the instrument was 1/350 while with the wider slit, the resolution was 1/200. The lower resolution is permissible in the latter case since the closest ion current is of mass 223 ($\text{Te}^{128}\text{F}_5^+$). The spectrum in Figure 6 was traced using the narrow collector slit.

The $\text{Te}^{130}/\text{Te}^{122}$ ratio is obtained directly by means of a five-figure integrating digital voltmeter-ratiometer and a digital printer in the measuring circuit of the mass spectrometer. The $\text{Te}^{122}\text{F}_5^+$ and $\text{Te}^{130}\text{F}_5^+$ ion beams are collected simultaneously and are amplified by vibrating reed electrometers. The currents are converted to voltages by means of 10^{11} and 10^{10} ohm grid leak resistors for the 122 and 130 mass currents respectively. The integrating digital voltmeter integrates an input voltage over a selected time and converts it to a proportional frequency. In passing through a counter section, this frequency is referred to a time base frequency so that the number printed is really a ratio of the converted frequency to the time base frequency. If an additional voltage-to-frequency converter is used, a second voltage can be converted into a frequency and used

in place of the time base frequency in the IDVM. The IDVM then is a ratiometer and displays the ratio of the two voltages.

The $\text{Te}^{130}/\text{Te}^{122}$ ratios of a standard sample and an unknown sample can be calculated directly from the printed output providing the major ion currents of the two samples are matched. Because of memory effects, it was necessary to allow at least five minutes after switching of the magnetic valves before recording the ratio in order to get reproduceable results. Approximately six sets each of ten printed ratios for each sample were obtained.

TABLE VII

Ion	% Abundance
TeF_5^+	58.3
TeF_4^+	2.8
TeF_3^+	5.5
TeF_2^+	2.8
TeF^+	0.4
Te^+	6.5
TeF_4^{++}	4.2
TeF_3^{++}	7.9
TeF_2^{++}	4.6
TeF^{++}	3.6
Te^{++}	3.3

RESULTS AND DISCUSSION

The results of the theoretical calculations as summarized in Tables IV and VI indicate that the $\text{Te}^{130}/\text{Te}^{122}$ ratio may be altered by over 3% in hypothetical reactions.

The experimental observations are consistent with some of the theoretical predictions. The measured $\text{Te}^{130}/\text{Te}^{122}$ ratios for the samples analyzed were referred to the isotopic composition of a standard sample of telluride from Copper Cliff, Ontario. The isotopic deviation from the standard is expressed in δ_{130} units where

$$\delta_{130} = \left[\frac{(\text{Te}^{130}/\text{Te}^{122})_{\text{sample}}}{(\text{Te}^{130}/\text{Te}^{122})_{\text{standard}}} - 1 \right] \times 1000$$

Generally speaking, a standard deviation of 0.02 percent could be realized in the measurements. The results are summarized in Table VIII and depicted graphically in Fig. 7.

A. Chemical reduction of potassium tellurite

Analysis showed that the tellurium precipitated in a 10 percent reduction of tellurite was 0.71 percent depleted in the heavier isotope in comparison with the initial tellurite. Using equation (19) developed in the theory, this value can be extrapolated with the assumption of first order kinetics to give the ratio of the rate constants k_{122}/k_{130} for the competitive reactions

THE UNIVERSITY OF CHICAGO
LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY

THE UNIVERSITY OF CHICAGO LIBRARY
1207 EAST 58TH STREET, CHICAGO, ILL. 60637
TEL: (312) 937-1234 FAX: (312) 937-1235
WWW.CHICAGO.LIBRARY.EDU
The University of Chicago Library is a member of the Association of Research Libraries (ARL) and the Association of American Universities (AAU). It is also a member of the American Library Association (ALA) and the American Association of University Libraries (AAUL).

CHICAGO LIBRARY

CHICAGO LIBRARY
1207 EAST 58TH STREET, CHICAGO, ILL. 60637
TEL: (312) 937-1234 FAX: (312) 937-1235
WWW.CHICAGO.LIBRARY.EDU

CHICAGO LIBRARY
1207 EAST 58TH STREET, CHICAGO, ILL. 60637
TEL: (312) 937-1234 FAX: (312) 937-1235
WWW.CHICAGO.LIBRARY.EDU

CHICAGO LIBRARY

CHICAGO LIBRARY


CHICAGO LIBRARY
1207 EAST 58TH STREET, CHICAGO, ILL. 60637
TEL: (312) 937-1234 FAX: (312) 937-1235
WWW.CHICAGO.LIBRARY.EDU

CHICAGO LIBRARY

Table VIII $\text{Te}^{130}/\text{Te}^{122}$ FRACTIONATION STUDY

Sample No.	Description	δ^{130}
1.	Commercial potassium tellurite B. D. H. Lot #54135	0.0 ± 0.2
2.	Elemental tellurium product of 10 percent chemical reduction of tellurite of sample 1	-7.1 ± 0.3
3.	Elemental tellurium, 41 to 53 percent reduction fraction of microbiological reduction of tellurite of sample 1	-2.1 ± 0.2
4.	Cripple Creek, Colorado (Sylvanite)	0.0 ± 0.2
5.	Copper Cliff, Ontario	0.0 ± 0.3
6.	Okanagan, B. C., PreCambrian Mines, See reference 137. (Tetradymite)	-4.0 ± 0.5

FIGURE 7 $\text{Te}^{122/130}$ FRACTIONATION STUDY





1000

1000


1000

1000

1000

1000

1000



This extrapolation gives $k_{122}/k_{130} = 1.0071 \pm 0.0003$ meaning that the lighter isotopic ion ruptures 0.7 percent faster. This is to be compared with the theoretically predicted results of Table VI. Looking at Table VI, "most reasonable" values of 1.00568 and 1.00718 for mass and atom fragments respectively are predicted. TeO_3 has a pyramidal XY_3 structure as shown in Figure 12, Appendix II. The bond being broken is equally as strong as the other bonds so that by Wolfsberg's reasoning (136), the reduced mass term will be closer to the reduced mass of the Te - O diatomic bond. Thus the value of 1.0071 ± 0.0003 compares very closely to the predicted result of 1.00718 for the atom-fragment model. This is also consistent with results found for selenite reduction (89).

B. Microbiological reduction study


For measurement, the elemental tellurium fraction precipitated between 41 and 53 percent reduction was used. This value can also be extrapolated to give the ratio k_{122}/k_{130} . The extrapolation is however more difficult and is shown graphically in Figure 8. The quantity "r" in equation (19) is for the integrated isotopic composition of the product.

THE HISTORY OF THE CITY OF LONDON

The history of the city of London is a subject of great interest and importance. It is a city of great antiquity, and its history is full of interest and variety. The city has been the seat of power and wealth for many centuries, and its history is a record of the growth and development of the British Empire. The city has been the center of commerce and industry, and its history is a record of the progress of the human race. The city has been the seat of learning and culture, and its history is a record of the advancement of the human mind. The city has been the center of the world, and its history is a record of the history of the world.

The history of the city of London is a subject of great interest and importance. It is a city of great antiquity, and its history is full of interest and variety. The city has been the seat of power and wealth for many centuries, and its history is a record of the growth and development of the British Empire. The city has been the center of commerce and industry, and its history is a record of the progress of the human race. The city has been the seat of learning and culture, and its history is a record of the advancement of the human mind. The city has been the center of the world, and its history is a record of the history of the world.

FIGURE 8 VARIATION OF ISOTOPIC COMPOSITION
OF PRODUCT WITH PERCENT REACTION



1. The first part of the paper is devoted to a study of the properties of the function $f(x)$ defined by the equation

$$f(x) = \int_0^x \frac{1}{1+t^2} dt$$



Three plots of "r" as a function of percentage reaction are shown in Figure 8 as dashed lines for values of $k_{122}/k_{130} = 1.005, 1.006, \text{ and } 1.007$. The point X indicates that if a reaction whose $k_{122}/k_{130} = 1.007$ has gone 40 percent to completion, the $\text{Te}^{122}/\text{Te}^{130}$ ratio of that total 40 percent is 1.0053 times that of the initial reactant.

For the microbiological study, however, a plot of isotopic composition of fractions vs. percentage reaction is necessary. Five plots for $k_{122}/k_{130} = 1.0050, 1.0055, 1.0060, 1.0065, \text{ and } 1.0070$ are shown by solid lines in Fig. 8. The information given by such a plot is represented by construction Y. If $k_{122}/k_{130} = 1.007$, and it is desired to know the average isotopic composition for the fraction between 10 percent and 20 percent reduction, the horizontal line is drawn at Y so that the shaded areas above and below the line are equal. In this example, the $\text{Te}^{122}/\text{Te}^{130}$ ratio of the 10 to 20 percent fraction of the product is 1.0058 times that of the initial reactant.

The above construction is performed in reverse for the experimental result. At Z, the horizontal line is the mean value of

$$\frac{(\text{Te}^{122}/\text{Te}^{130})_{\text{reduced fraction}}}{(\text{Te}^{122}/\text{Te}^{130})_{\text{initial reactant}}} = 1.0021$$

The first of these is the fact that the system is not a simple one, but a complex one, involving many different factors.

The second is the fact that the system is not a static one, but a dynamic one, involving many different factors.

The third is the fact that the system is not a homogeneous one, but a heterogeneous one, involving many different factors.

The fourth is the fact that the system is not a uniform one, but a non-uniform one, involving many different factors.

The fifth is the fact that the system is not a simple one, but a complex one, involving many different factors.

The sixth is the fact that the system is not a static one, but a dynamic one, involving many different factors.

The seventh is the fact that the system is not a homogeneous one, but a heterogeneous one, involving many different factors.

The eighth is the fact that the system is not a uniform one, but a non-uniform one, involving many different factors.

The ninth is the fact that the system is not a simple one, but a complex one, involving many different factors.

The tenth is the fact that the system is not a static one, but a dynamic one, involving many different factors.

The eleventh is the fact that the system is not a homogeneous one, but a heterogeneous one, involving many different factors.

The twelfth is the fact that the system is not a uniform one, but a non-uniform one, involving many different factors.

The thirteenth is the fact that the system is not a simple one, but a complex one, involving many different factors.

for the fraction between 41 and 53 percent reaction. It is seen that the line for $k_{122}/k_{130} = 1.0060$ very closely equalizes the shaded areas. The measurement error of ± 0.02 percent is represented by the vertical line at Z and its extremities touch the curves corresponding to $k_{122}/k_{130} = 1.0055$ and 1.0065 . Therefore, for the bacterial reduction, $k_{122}/k_{130} = 1.0060 \pm 0.0005$. This compares very favourably with the result of the chemical reduction study and suggests that in the bacterial reduction, the effect noted arises essentially in tellurium-oxygen bond breakage.

k_{122}/k_{130} can be determined more accurately by measurements made of fractions produced near the beginning of the reaction. Such samples are available, but measurements are being withheld until the mass spectrometer is adapted to handle smaller samples.

C. Natural isotopic abundance studies

Two of the natural samples and the commercial tellurite have the same $\text{Te}^{122}/\text{Te}^{130}$ ratio within the measurement errors. Similarly in the survey of natural selenium isotopes (60) a large number of samples had a similar $\text{Se}^{76}/\text{Se}^{82}$ ratio. In both cases, the constancy of isotopic composition can be related to high temperature processes where the isotope fractionation is very small. (The isotope fractionation generally decreases with temperature as exemplified in the calculations

The first of these is the fact that the system is not a simple one. It is a complex system, and the behavior of the system is not linear. The system is a complex system, and the behavior of the system is not linear. The system is a complex system, and the behavior of the system is not linear.

The second of these is the fact that the system is not a simple one. It is a complex system, and the behavior of the system is not linear. The system is a complex system, and the behavior of the system is not linear. The system is a complex system, and the behavior of the system is not linear.

The third of these is the fact that the system is not a simple one. It is a complex system, and the behavior of the system is not linear. The system is a complex system, and the behavior of the system is not linear. The system is a complex system, and the behavior of the system is not linear.

The System is a Complex System

The system is a complex system, and the behavior of the system is not linear. The system is a complex system, and the behavior of the system is not linear. The system is a complex system, and the behavior of the system is not linear. The system is a complex system, and the behavior of the system is not linear. The system is a complex system, and the behavior of the system is not linear.

of Tables IV and VI.) In the selenium study (60), large alterations to the $\text{Se}^{76}/\text{Se}^{82}$ ratio occurred only in soil and plant samples where the temperature of the processes would be very low. In the tellurium study, however, the sample of tetradymite from the west shore of the Okanagan Lake was 0.4 percent depleted in Te^{130} . This result is surprising. A measurement of yield rules out any possibility that the sample was isotopically altered in the chemical extraction. This sample behaved the same as the others in hexafluoride preparation and in mass spectrometric analysis. It would seem that the result is real but it should be interpreted as one measurement of one sample. No attempt can be made to explain this result until more natural samples are examined.

In conclusion, therefore, this preliminary survey shows that the $\text{Te}^{122}/\text{Te}^{130}$ ratio can be altered chemically and microbiologically in agreement with theoretical predictions. Further, it would seem that a more extensive study of natural samples might be beneficial in elucidating natural processes.

The first part of the paper discusses the importance of the
theoretical framework in the study of the
relationship between the variables. The second part
presents the empirical results of the study. The third part
discusses the implications of the findings for the
theory and practice. The fourth part concludes the paper.

The results of the study show that there is a significant
relationship between the variables. The findings
support the theoretical framework. The implications
for the theory and practice are discussed.

The study has some limitations. Further research is
needed to explore the relationship between the variables
in more detail. The findings of the study are
generalizable to other contexts.

BIBLIOGRAPHY

1. Aldrich, L. T., and Nier, A. O., Phys. Rev., 74, 876, 159, (1948)
2. Aston, F. W., Phil. Mag. 38, 707, (1919); 39, 449, (1920)
3. Aston, F. W., Phil. Mag. 49, 1191, (1925)
4. Aston, F. W., Proc. Roy. Soc. (London), A132, 487, (1931)
5. Bach F, Bonhoeffer, K. F., and Fajans, E., Zeits. für Physik chemie, A168, 313, (1934)
6. Baertschi, P., Nature 168, 288, (1951)
7. Bainbridge, K. T., Phys. Rev. 39, 1021, (1932)
8. Bartlett, James H. Jr., Phys. Rev. 45, 847, (1934)
9. Bigeleisen, J., J. Chem. Phys. 17, 675 (1940)
10. Bigeleisen, J., and Mayer, M.G., J. Chem. Phys., 15, 261, (1947)
11. Bigeleisen, J., J. Chem. Phys., 17, 675, (1949)
12. Bigeleisen, H., and Friedman, L., J. Chem. Phys., 17, 998, (1949)
13. Bird, Marjorie L., and Challenger, Frederick, J. Chem. Soc. p163, (1939)
14. Bose, H., and Ray, B. B., Indian J. Phys., 15, 233, (1941)
15. Butlin, K. R., Research 6, 184, (1953)
16. Carr, F., and Pearson, T. G., J. Chem. Soc., p 282, (1938)

17. Challenger, F., Higginbottom, C., and Ellis, L., J. Chem. Soc., page 95 (1933)
18. Ch'in, T. L., Proc. Soc. Expt'l. Med., 38, 700, (1938)
19. Clarke, F. W., and Washington, H. S., Proc. Nat. Acad., 8, 108, (1952)
20. Clayton, R. N., J. Chem. Phys., 30, 1246, (1955)
21. Cohen, E., and Kröner, J. F., Zeit. Phys. Chem., 82, 587, (1913)
22. Craig, H., Geochim. et Cosmochim. Acta., 3, 53, (1953)
23. Craig, H., and Boato, G., Am. Rev. Phys. Chem., 6, 403, (1955)
24. Davy, H., Phil. Trans., 100, 161, (1810)
25. Dempster, A. J., Phys. Rev., 11, 316, (1918)
26. Dempster, A. J., Phys. Rev., 50, 186, (1936)
27. Dodwell, L. G., Proc. Ind. Acad. Sci., 55, (1945)
28. Duckworth, H. E., Kegley, C. L., Olson, J. M., and Stanford, G. S., Phys. Rev. 83, 1114, (1951)
29. DuFresne, Ann., Geochim et Cosmochim Acta, 20, 141, (1960)
30. Duval, Clement, "Inorganic Thermo-gravimetric Analysis", pp. 302, 384, Elsevier, New York, (1953)
31. Eidinoff, M. L., Trans. N. Y. Acad. Sci., 16, 76, (1953)
32. Epstein, S., Buchsbaum, R., Lowenstem, H., and Urey, H. C., Bull. Geol. Soc. Amer., 62, 417, (1953)

33. Epstein, S., "Mass Spectroscopy in Physical Research", National Bur. Std. U. S. Circ. 522, page 133, (1953)
34. Epstein, S., "Oxygen Isotope Measurements in Glacial Ice", Nat. Acad. Sci. - Nat. Res. Council Bull. 845, p. 1521, (1960)
35. Evans, M. G., and Polanyi, M., Trans. Far. Soc., 31, 875, (1935)
36. Eyring, H., J. Chem. Phys., 3, 107, (1935)
37. Farkas, L., and Farkas, A., Naturwiss, 22, 218, (1934)
38. Freely, H. W., and Kulp, J. L., Bull. Amer. Assoc. Petrol. Geol., 41, 1802, (1957)
39. Friedman, I., Geochim. et Cosmochim Acta, 4, 89, (1953)
40. Gaunt, J., Trans. Far. Soc., 50, 546, (1954)
- 40a. Glasstone, S., Laidler, K. J., and Eyring, H., "The Theory of Rate Processes", McGraw Hill, N. Y., (1941)
41. Glockler, G., and Tung, J. Y., J. Chem. Phys., 13, 338, (1945)
42. Gooch, F. A., and Peirce, A. W., Am. Jour. Sci., 1, 181, (1896)
43. Halsted, Richard E., Phys. Rev. 88, 666, (1952)
44. Harrison, A., and Thode, H. G., Trans. Far. Soc., 54, 84, (1958)
45. Heath, D. F., and Linnett, J. W., Trans. Far. Soc., 44, 878, (1948)
46. Heath, D. F., and Linnett, J. W., Trans. Far. Soc., 45, 264, (1949)

THE
JOURNAL
OF
THE
ROYAL ANTHROPOLOGICAL INSTITUTE
OF GREAT BRITAIN AND IRELAND
VOLUME 10
PART 1
1880
LONDON
PUBLISHED BY THE INSTITUTE
21, BEDFORD SQUARE, W.C.

47. Heslop, R. B., and Robinson, P. L., "Inorganic Chemistry", Elsevier Publishing Company, New York, p 352, (1963)
48. Herzberg, G., "Spectra of Diatomic Molecules", D. Van Nostrand, Princeton, New Jersey, (1950)
49. Herzberg, G., "Infrared and Raman Spectra", D. Van Nostrand Co. Inc., Princeton, New Jersey, (1956)
50. Hill, R. A., Edwards, T. H., Rossman, K., Harahari Rao, K., Nielsen, H. H., J. Mol. Spectry., 14(3), 221, (1964)
51. Hillebrand, W. F., Lundell, G. E. R., Bright, H. A., Hoffman, J. I., "Applied Inorganic Analysis", pp. 228, 334, Wiley, New York, (1953)
52. Hirschfelder, J. O., and Wigner, J., J. Chem. Phys., 1, 616, (1939)
53. Hoering, T. C., Science, 122, 1233, (1955)
54. Hollander, J. M., Perlman, I., and Seaborg, G. T., Revs. Modern Phys., 25, 469, (1953)
55. Johnson, C. W., and Krouse, R. H., Unpublished work, University of Alberta, Edmonton, Alberta, (1964)
56. Junkins, J. H. Bernhardt, H. A., and Barber, E. J., J. Am. Chem. Soc., 74, 5749, (1952)
57. Kaplan, I. R., and Rittenberg, S. C., J. gen. Microbiol, 34, 195, (1964)
58. Keller, Edward, J. Am. Chem. Soc., 19, 771, (1897)
59. King, W. E., and Davis, L., Am. Jour. of Pub. Health, 4, 917, (1914)
60. Krouse, R. H., and Thode, H. G., Can. Jour. Chem., 40, 367, (1958)

1875

1876

1877

1878

1879

1880

1881

1882

1883

1884

1885

61. Kruse, P. W., McGlauchin, L. D., and McQuisten, R. B., "Elements of Infrared Technology", John Wiley and Sons, Inc., New York, (1962)
62. Landergren, V. S., Z. Naturf., 8b, 537, (1953)
63. Lindsay, J. G., Bourns, A. N., and Thode, H. G., Can. J. Chem., 29, 192, (1951)
64. Lindsay, J. G., Bourns, A. N., and Thode, H. G., Can. J. Chem., 30, 163, (1952)
65. Lindsay, J. G., McElheran, D. E., and Thode, H. G., Journ. Chem. Phys., 17, 589, (1949)
66. Linnett, J. W., and Simpson, C. J. S. M., Trans. Far. Soc., 55, 857, (1959)
67. Lippincott, E. R., and Dayhoff, M. O., Spectrochim Acta, 16, 807, (1960)
68. Macnamara, J., and Thode, H. G., Phys. Rev., 78, 307, (1950)
69. Macnamara, J., and Thode, H. G., Research 4, 582, (1951)
70. Macnamara, J., Fleming, W., Szabo, A., and Thode, H. G., Can. J. Chem., 30, 73, (1952)
71. Mars, K. E., Jour. Geol., 59, 131, (1951)
72. Mayer, J. E., and Mayer, M. G., "Statistical Mechanics", p. 207 ff., J. Wiley and Sons, Inc., N. Y., (1940)
73. McCrea, J. M., J. Chem. Phys., 18, 849, (1950)
74. McElheran, P. E., M. Sc. Thesis, McMaster University, Hamilton, Ontario, (1951)
75. McKinney, C. R., McCrea, J. M., Epstein, S., Allen, H. H., and Urey, H. C., Rev. Sci. Inst., 21, 724, (1950)

1. The first part of the paper is devoted to a general discussion of the problem.

2. In the second part, we consider the case of a single particle.

3. The third part is devoted to the case of a system of particles.

4. In the fourth part, we consider the case of a continuous medium.

5. The fifth part is devoted to the case of a system of continuous media.

6. In the sixth part, we consider the case of a single continuous medium.

7. The seventh part is devoted to the case of a system of continuous media.

8. In the eighth part, we consider the case of a single continuous medium.

9. The ninth part is devoted to the case of a system of continuous media.

10. In the tenth part, we consider the case of a single continuous medium.

11. The eleventh part is devoted to the case of a system of continuous media.

76. Mellor, J. W., "A Comprehensive Treatise on Inorganic and Theoretical Chemistry", p. 5, Longman's Green and Co., Toronto, (1931)
77. Migeotte, R., Bull. Soc. Roy. Sci..Liege, 11, 44, (1942)
78. Miller, L. P., Contrib. Boyce Thompson Inst., 15, 437, (1949)
79. Morton, Harry E., and Anderson, Thomas F., Proc. Soc. Expt'l Biol. Med., 46, 272, (1941)
80. Munther, V., and Dansgard, W., Medd. Dansk. Geol. Foren, 12, 159, (1951)
81. Nakai, N., and Jensen, M. L., Geochim et Cosmochim Acta, 28, 1893, (1964)
82. Nakomoto, K., "Infrared Spectra of Inorganic and Co-ordination Compounds", John Wiley and Sons, Inc., New York, (1963)
83. Nier, A. O., and Gulbransen, E. J., J. Am. Chem. Soc., 61, 697, (1939)
84. Nier, A. O., and Murphy, B. F., Phys. Rev., 59, 771, (1941)
85. Pistorius, C. W. F. T., J. Chem. Phys., 27, 965, (1957)
86. Rankama, K., Bull. Geol. Soc. Amer., 59, 389, (1948)
87. Redlich, O., Z. Physik. Chem., B28, 371, (1935)
88. Reed, James. F., Analytical Chem., 32, 662, (1960)
89. Rees, C. E. Unpublished Ph. D. Thesis, McMaster University, Hamilton, Ontario, (1964)
90. Richards, T. W., and Lembergt, M. C., J. Am. Chem. Soc., 36, 1329, (1914)

The first part of the paper discusses the importance of the
second part of the paper discusses the importance of the
third part of the paper discusses the importance of the
fourth part of the paper discusses the importance of the
fifth part of the paper discusses the importance of the
sixth part of the paper discusses the importance of the
seventh part of the paper discusses the importance of the
eighth part of the paper discusses the importance of the
ninth part of the paper discusses the importance of the
tenth part of the paper discusses the importance of the

91. Rittenberg, D., Bleakney, W., and Urey, H. C.,
J. Chem. Phys., 2, 98, (1934)
92. Ropp, G. A., Nucleonics, 10, #10, 22, (1952)
93. Rossmann, Kurt, and Straley, Joseph W., J. Chem.
Phys., 24, 1276, (1956)
94. Semat, Henry, "Introduction to Atomic and Nuclear
Physics", Rinehart and Co. Inc., New
York, (1960)
95. Siebert, H., Z. anorg. u. allgem. chem., 275,
225, (1955)
96. Sienko, M. J., and Plane, R. A., "Chemistry",
McGraw-Hill Book Co. Inc., New York,
(1957)
97. Silverman, S. R., and Epstein, S., Bull. Amer.
Ass. Petrol. Geol., 42, 998, (1958)
98. Sindeeva, N. D., "Mineralogy and Types of Deposits
of Selenium and Tellurium", Interscience
Publishers, New York, p. 255, (1964)
99. Smith, S., and Linnett, J. W., Trans. Far. Soc.,
52, 891, (1956)
100. Soddy, F., Chem. Soc. Ann., Rep. 1, 285, (1910)
101. Stevenson, D. P., Jour. Chem. Phys., 8, 285, (1940)
102. Sula, L., Am. Rev. Tuberc., 56, 241, (1947)
103. Thanalakshmi, Miss R., Current Science, 31, 374,
(1962)
104. Thode, H. G. Macnamara, J., and Collins, C. B.,
Can. J. Research, 27B, 361, (1947)
105. Thode, H. G., Kleerekoper, H., and McElheran,
D. E., Research, 4, 581, (1951)
106. Thode, H. G., Ann. Rev. Phys., Chem., 4, 95, (1953)

107. Thode, H. G., Wanless, R. K., and Wallaach, R.,
Geochim et Cosmochim Acta, 5, 286, (1954)
108. Thode, H. G., Bulletin of Canadian Petroleum
Geology, 12, 246, (1963)
109. Thomson, J. J., "Rays of Positive Electricity and
their Applications to Chemical Analysis",
Longman's Green and Company, London,
(1913)
110. Trofimov, A., Doklady Akad. Nauk., SSR85, 169,
(1952)
111. Trudge, A. P., and Thode, H. G., "Thermodynamic
Properties of Isotopic Compounds of
Sulphur", Can. Jour. Res., 28B, 567, (1950)
112. Urey, H. C., Brickwedde, F. G., and Murphy, G. M.,
Phys. Rev., 39, 164, 864, (1932)
113. Urey, H. C., and Washburn, E. W., Proc. Nat.
Acad. Sci., 18, 496, (1932)
114. Urey, H. C., and Rittenberg, D. J., Jour. Chem.
Phys., 1, 137, (1933)
115. Urey, H. C., and Rittenberg, D. J., J. Am. Chem.
Soc., 56, 1885, (1934)
116. Urey, H. C., and Greiff, L., J. Am. Chem. Soc.,
57, 321, (1935)
117. Urey, H. C., and Teal, G. K., Rev. Mod. Phys.,
1, 34, (1935)
118. Urey, H. C., J. Chem. Soc., p562, (1947)
119. Urey, H. C., Science, 108, 489, (1948)
120. Urey, H. C., Epstein, S., McKinney, C., and McCrea,
J. M., Bull. Geol. Soc. Amer., 59,
1359, (1948)
121. Venkateswarlu, K., Somasundaram, V., and Krishna,
M. G., Z. Physik, Chem., 212, 145, (1960)

- 122. Vernon, Richard Henry, J. Chem. Soc., p. 86,
(1920)
- 123. Vinogradov, A. P., "The Geochemistry of Rare and
Dispersed Chemical Elements in the Soil",
trans. from Russian-Consultants' Bureau,
N.Y., (1959)
- 124. Voge, H. H., and Libby, W. F., J. Am. Chem.
Soc., 59, 2474, (1937)
- 125. Wachstein, M., Proc. Soc. Exptl. Biol. Med.,
72, 175, (1949)
- 126. Wanless, R. K., and Thode, H. G., J. Sci. Inst.,
30, 395, (1953)
- 127. Weber, L. A., Wahl, M. H., and Urey, H. C., J.
Chem. Phys., 3, 129, (1935)
- 128. Wells, A. F., and Bailey, M., J. Chem. Soc., p.1282,
(1949)
- 129. Wentink, Tunis Jr., J. Chem. Phys., 30, 105, (1959)
- 130. West, S. S., Geophysics, 10, 406, (1945)
- 131. White, J. R., and Cameron, A. F., Phys. Rev.,
74, 991, (1948)
- 132. Williams, D., and Yuster, P., Phys. Rev., 69,
556, (1946)
- 133. Wickman, F. E., and von Ubish, H., Geochim et
Cosmochim Acta, 1, 119, (1951)
- 134. Wickman, F. E., Geochim et Cosmochim Acta, 2, 243,
(1952)
- 135. Wickman, F. E., Nature, 169, 1051, (1951)
- 136. Wolfsberg, M., J. Chem. Phys., 33, 21, (1960)
- 137. Summary report, Geological Survey of Canada, Part b,
"White Elephant Mines - PreCambrian Mines",
(1931)

The first part of the paper is devoted to a general discussion of the problem. It is shown that the problem is of great importance in the theory of differential equations. The second part is devoted to the study of the properties of the solutions of the equation. It is shown that the solutions of the equation are unique and that they depend continuously on the initial conditions. The third part is devoted to the study of the asymptotic properties of the solutions. It is shown that the solutions of the equation tend to zero as $t \rightarrow \infty$. The fourth part is devoted to the study of the stability of the solutions. It is shown that the solutions of the equation are stable. The fifth part is devoted to the study of the periodic properties of the solutions. It is shown that the solutions of the equation are periodic. The sixth part is devoted to the study of the ergodic properties of the solutions. It is shown that the solutions of the equation are ergodic. The seventh part is devoted to the study of the mixing properties of the solutions. It is shown that the solutions of the equation are mixing. The eighth part is devoted to the study of the entropy properties of the solutions. It is shown that the solutions of the equation have a positive entropy. The ninth part is devoted to the study of the topological properties of the solutions. It is shown that the solutions of the equation are topologically transitive. The tenth part is devoted to the study of the dynamical properties of the solutions. It is shown that the solutions of the equation are dynamical systems.

APPENDIX I

CALCULATIONS OF ISOTOPIC VIBRATIONAL FREQUENCY SHIFTS
FOR TELLURIUM - CONTAINING MOLECULES

Values for the normal vibrational frequencies of many tellurium-containing molecules are available from infra-red spectra and can be found in various publications. However, since no consideration is given for differing isotopic masses therein, it is assumed that these experimental values apply to the most abundant tellurium isotope, Te^{130} . The frequencies of the Te^{122} containing molecules were then calculated from these known frequencies using normal force equations which describe molecules of different symmetries.

A. Diatomic Molecules

In considering diatomic molecules, the anharmonic oscillator is chosen as a model. The potential energy function is then given in the form

$$U = f(r - r_e)^2 - g(r - r_e)^3 + . . . \quad (\text{I.1})$$

where r_e is the equilibrium internuclear distance and $(r - r_e)$ is the displacement from this equilibrium position.

Schroedinger's equation for such a model then becomes

$$\nabla^2 \psi + \frac{8\pi^2 \mu}{h^2} (E - U) \psi = 0 \quad (\text{I.2})$$

If the anharmonicity is small, ($g \ll f$), the energy values of this equation will be the eigenvalues given by

$$E_{\text{vib}} = hc \left[\omega_e \left(v + \frac{1}{2}\right) - \omega_e x_e \left(v + \frac{1}{2}\right)^2 + \omega_e y_e \left(v + \frac{1}{2}\right)^3 \dots \right] \quad (\text{I.3})$$

where v is the vibrational quantum number, ω_e is the classical vibrational frequency in cm^{-1} , and μ is the reduced mass of the two atoms, $\left(\frac{m_1 m_2}{m_1 + m_2}\right)$.

Classically, for a harmonic oscillator

$$\nu_{\text{osc}} = \frac{1}{2\pi} \sqrt{\frac{K}{\mu}} \quad \text{sec}^{-1}$$

where ν_{osc} is the vibrational frequency and the restoring force is $-K(r - r_e)$ during a displacement of $(r - r_e)$.

The potential energy curves for isotopic molecules are identical to a high degree since they depend only on the motions of the electrons and the coulomb repulsion of the nuclei (48). This means that the force constant K can

be assumed independent of isotopic substitution. Therefore the ratio of the classical frequencies between two isotopic molecules would be given as

$$\frac{\nu_1}{\nu_2} = \sqrt{\frac{\mu_2}{\mu_1}}$$

Or in terms of the most abundant isotope,

$$\frac{\nu_{\text{isotope}}}{\nu} = \sqrt{\frac{\mu}{\mu_{\text{isotope}}}} = \rho$$

From the relationship that $\nu = c\omega$,

$$\frac{\omega_e}{\omega} = \rho \quad (\text{I.4})$$

Thus from equations I.3 and I.4 the energy levels of the isotopic molecule are given by

$$(E_v)_i = hc \left[\rho \omega_e \left(v + \frac{1}{2}\right) - \rho^2 \omega_e x_e \left(v + \frac{1}{2}\right)^2 + \dots \right] \quad (\text{I.5})$$

The only vibrational energy transitions possible are those dictated by the selection rule, i.e., $\Delta v = \pm 1$. Thus the vibrational energy change in a transition from $v = 0$ to $v = 1$ is given by equation I.3 as

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
530 SOUTH EAST ASIAN AVENUE
CHICAGO, ILLINOIS 60607-7070

7/1/88

Dear Professor [Name]:
I am writing to you regarding the [Topic].

I have been thinking about the [Topic] and the [Topic].

I am looking forward to the [Topic] and the [Topic].

I am looking forward to the [Topic] and the [Topic].

I am looking forward to the [Topic] and the [Topic].

$$\begin{aligned}\Delta E_v &= hc [\omega_e - 2\omega_e x_e + \dots] \\ &= hc\omega\end{aligned}\quad (I.6)$$

From I.4, the energy change for a $v = 1$ to $v = 0$ transition for the isotopic molecule is given in terms of the transition for the abundant molecule as

$$\begin{aligned}(\Delta E_v)_i &= hc [\rho\omega_e - 2\rho^2\omega_e x_e + \dots] \\ &= hc\omega_i\end{aligned}\quad (I.7)$$

From equation (I.7), the expression for the vibrational frequency of the isotopic molecule in terms of the frequency of the more abundant molecule as

$$\omega_i = \rho\omega_e - 2\rho^2\omega_e x_e + \dots \quad (I.8)$$

Experimental values for ω_e and $\omega_e x_e$ for the Te^{130} -containing molecules are obtained from observed data (48), (67). Then from I.4,

$$\omega_e^{122} = \sqrt{\frac{\mu^{130}}{\mu^{122}}} \cdot \omega_e^{130} = \rho\omega_e^{130}$$

and

$$(\omega_e x_e)^{122} = (\rho)^2 (\omega_e x_e)^{130} .$$

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO
DIVISION OF THE PHYSICAL SCIENCES
DEPARTMENT OF CHEMISTRY

1954-1955

THE UNIVERSITY OF CHICAGO
DIVISION OF THE PHYSICAL SCIENCES
DEPARTMENT OF CHEMISTRY

THE UNIVERSITY OF CHICAGO
DIVISION OF THE PHYSICAL SCIENCES
DEPARTMENT OF CHEMISTRY

THE UNIVERSITY OF CHICAGO
DIVISION OF THE PHYSICAL SCIENCES
DEPARTMENT OF CHEMISTRY

THE UNIVERSITY OF CHICAGO
DIVISION OF THE PHYSICAL SCIENCES
DEPARTMENT OF CHEMISTRY

Table IX gives the calculations for the vibrational frequencies of Te^{122} -containing molecules from the available data. These were computed with the IBM 7040 computer program found on page A37, Appendix II.

B. Polyatomic Molecules

In these molecules, a simple model cannot be chosen for the vibrational frequencies. Detailed information must be obtained about the structure of the molecule in order to describe the force fields and set up equations for the vibrational frequencies. The force equations which seemed to best describe the observed spectra were chosen. The force constants were solved in terms of the most abundant Te^{130} isotope and experimentally observed frequencies. Then the other isotopic masses were substituted to find the corresponding isotopic frequencies.

In several cases even the best description of the molecule did not give results in perfect agreement with the observed vibrational frequencies. Therefore, linear correction factors are used when needed to align the calculated frequencies to those observed.

THE UNIVERSITY OF CHICAGO

PH.D. THESIS
SUBMITTED TO THE FACULTY OF THE DIVISION OF THE PHYSICAL SCIENCES
IN CANDIDACY FOR THE DEGREE OF DOCTOR OF PHILOSOPHY
DEPARTMENT OF CHEMISTRY

BY

JOHN EDWARD HARRIS
B.A. 1961, M.A. 1963, Ph.D. 1965
THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
5408 S. UNIVERSITY AVE.
CHICAGO, ILL. 60637
1965

THE UNIVERSITY OF CHICAGO
LIBRARY
5408 S. UNIVERSITY AVE.
CHICAGO, ILL. 60637

TABLE IX

Calculation of Vibrational Frequency Shifts for Te^{130} and Te^{122} -containing molecules

Molecule	ω_e^{130} cms ⁻¹	$(\omega_{e,x_e})^{130}$ cms ⁻¹	$\frac{\mu^{130}}{\mu^{122}}$	$\left[\frac{\mu^{130}}{\mu^{122}} \right]^{\frac{1}{2}}$	ω_e^{122} cms ⁻¹	ω_{e,x_e}^{122} cms ⁻¹	ω^{130} cms ⁻¹	ω^{122} cms ⁻¹
Ge^{74}Te	323.4	1.00	1.024	1.0118	327.2	1.024	321.40	325.18
Pb^{208}Te	211.8	0.12	1.040	1.0200	216.0	0.125	211.56	215.79
Sn^{120}Te	259.5	0.50	1.032	1.0156	263.6	0.516	258.50	262.52
TeO^{16}	796.0	3.50	1.007	1.0036	798.9	3.525	789.00	791.81
Si^{28}Te	480.4	1.16	1.012	1.0058	483.2	1.173	478.08	480.84
$\text{Te}^{130}\text{Te}^{130}$	251.5	0.55	1.033	1.0163	255.6	0.568	250.40	254.46
HTe	2250.0	41.30	1.001	1.0003	2250.6	41.32	2167.40	2167.40

17114	7024
17115	7025
17116	7026
17117	7027
17118	7028
17119	7029
17120	7030
17121	7031
17122	7032
17123	7033
17124	7034
17125	7035
17126	7036
17127	7037
17128	7038
17129	7039
17130	7040
17131	7041
17132	7042
17133	7043
17134	7044
17135	7045
17136	7046
17137	7047
17138	7048
17139	7049
17140	7050
17141	7051
17142	7052
17143	7053
17144	7054
17145	7055
17146	7056
17147	7057
17148	7058
17149	7059
17150	7060
17151	7061
17152	7062
17153	7063
17154	7064
17155	7065
17156	7066
17157	7067
17158	7068
17159	7069
17160	7070
17161	7071
17162	7072
17163	7073
17164	7074
17165	7075
17166	7076
17167	7077
17168	7078
17169	7079
17170	7080
17171	7081
17172	7082
17173	7083
17174	7084
17175	7085
17176	7086
17177	7087
17178	7088
17179	7089
17180	7090
17181	7091
17182	7092
17183	7093
17184	7094
17185	7095
17186	7096
17187	7097
17188	7098
17189	7099
17190	7100
17191	7101
17192	7102
17193	7103
17194	7104
17195	7105
17196	7106
17197	7107
17198	7108
17199	7109
17200	7110
17201	7111
17202	7112
17203	7113
17204	7114
17205	7115
17206	7116
17207	7117
17208	7118
17209	7119
17210	7120
17211	7121
17212	7122
17213	7123
17214	7124
17215	7125
17216	7126
17217	7127
17218	7128
17219	7129
17220	7130
17221	7131
17222	7132
17223	7133
17224	7134
17225	7135
17226	7136
17227	7137
17228	7138
17229	7139
17230	7140
17231	7141
17232	7142
17233	7143
17234	7144
17235	7145
17236	7146
17237	7147
17238	7148
17239	7149
17240	7150
17241	7151
17242	7152
17243	7153
17244	7154
17245	7155
17246	7156
17247	7157
17248	7158
17249	7159
17250	7160
17251	7161
17252	7162
17253	7163
17254	7164
17255	7165
17256	7166
17257	7167
17258	7168
17259	7169
17260	7170
17261	7171
17262	7172
17263	7173
17264	7174
17265	7175
17266	7176
17267	7177
17268	7178
17269	7179
17270	7180
17271	7181
17272	7182
17273	7183
17274	7184
17275	7185
17276	7186
17277	7187
17278	7188
17279	7189
17280	7190
17281	7191
17282	7192
17283	7193
17284	7194
17285	7195
17286	7196
17287	7197
17288	7198
17289	7199
17290	7200
17291	7201
17292	7202
17293	7203
17294	7204
17295	7205
17296	7206
17297	7207
17298	7208
17299	7209
17300	7210
17301	7211
17302	7212
17303	7213
17304	7214
17305	7215
17306	7216
17307	7217
17308	7218
17309	7219
17310	7220
17311	7221
17312	7222
17313	7223
17314	7224
17315	7225
17316	7226
17317	7227
17318	7228
17319	7229
17320	7230
17321	7231
17322	7232
17323	7233
17324	7234
17325	7235
17326	7236
17327	7237
17328	7238
17329	7239
17330	7240
17331	7241
17332	7242
17333	7243
17334	7244
17335	7245
17336	7246
17337	7247
17338	7248
17339	7249
17340	7250
17341	7251
17342	7252
17343	7253
17344	7254
17345	7255
17346	7256
17347	7257
17348	7258
17349	7259
17350	7260
17351	7261
17352	7262
17353	7263
17354	7264
17355	7265
17356	7266
17357	7267
17358	7268
17359	7269
17360	7270
17361	7271
17362	7272
17363	7273
17364	7274
17365	7275
17366	7276
17367	7277
17368	7278
17369	7279
17370	7280
17371	7281
17372	7282
17373	7283
17374	7284
17375	7285
17376	7286
17377	7287
17378	7288
17379	7289
17380	7290
17381	7291
17382	7292
17383	7293
17384	7294
17385	7295
17386	7296
17387	7297
17388	7298
17389	7299
17390	7300
17391	7301
17392	7302
17393	7303
17394	7304
17395	7305
17396	7306
17397	7307
17398	7308
17399	7309
17400	7310
17401	7311
17402	7312
17403	7313
17404	7314
17405	7315
17406	7316
17407	7317
17408	7318
17409	7319
17410	7320
17411	7321
17412	7322
17413	7323
17414	7324
17415	7325
17416	7326
17417	7327
17418	7328
17419	7329
17420	7330
17421	7331
17422	7332
17423	7333
17424	7334
17425	7335
17426	7336
17427	7337
17428	7338
17429	7339
17430	7340
17431	7341
17432	7342
17433	7343
17434	7344
17435	7345
17436	7346
17437	7347
17438	7348
17439	7349
17440	7350
17441	7351
17442	7352
17443	7353
17444	7354
17445	7355
17446	7356
17447	7357
17448	7358
17449	7359
17450	7360
17451	7361
17452	7362
17453	7363
17454	7364
17455	7365
17456	7366
17457	7367
17458	7368
17459	7369
17460	7370
17461	7371
17462	7372
17463	7373
17464	7374
17465	7375
17466	7376
17467	7377
17468	7378
17469	7379
17470	7380
17471	7381
17472	7382
17473	7383
17474	7384
17475	7385
17476	7386
17477	7387
17478	7388
17479	7389
17480	7390
17481	7391
17482	7392
17483	7393
17484	7394
17485	7395
17486	7396
17487	7397
17488	7398
17489	7399
17490	7400
17491	7401
17492	7402
17493	7403
17494	7404
17495	7405
17496	7406
17497	7407
17498	7408
17499	7409
17500	7410
17501	7411
17502	7412
17503	7413
17504	7414
17505	7415
17506	7416
17507	7417
17508	7418
17509	7419
17510	7420
17511	7421
17512	7422
17513	7423
17514	7424
17515	7425
17516	7426
17517	7427
17518	7428
17519	7429
17520	7430
17521	7431
17522	7432
17523	7433
17524	7434
17525	7435
17526	7436
17527	7437
17528	7438
17529	7439
17530	7440
17531	7441
17532	7442
17533	7443
17534	7444
17535	7445
17536	7446
17537	7447
17538	7448
17539	7449
17540	7450
17541	7451
17542	7452
17543	7453
17544	7454
17545	7455
17546	7456
17547	7457
17548	7458
17549	7459
17550	7460
17551	7461
17552	7462
17553	7463
17554	7464
17555	7465
17556	7466
17557	7467
17558	7468
17559	7469
17560	7470
17561	7471
17562	7472
17563	7473
17564	7474
17565	7475
17566	7476
17567	7477
17568	7478
17569	7479
17570	7480
17571	7481
17572	7482
17573	7483
17574	7484
17575	7485
17576	7486
17577	7487
17578	7488
17579	7489
17580	7490
17581	7491
17582	7492
17583	7493
17584	7494
17585	7495
17586	7496
17587	7497
17588	7498
17589	7499
17590	7500
17591	7501
17592	7502
17593	7503
17594	7504
17595	7505
17596	7506
17597	7507
17598	7508
17599	7509
17600	7510
17601	7511
17602	7512
17603	7513
17604	7514
17605	7515
17606	7516
17607	7517

TeF₆

Tellurium hexafluoride is of octahedral structure of O_h symmetry. The octahedral molecules, XY_6 , of symmetry O_h each show six fundamental vibration frequencies. Of these six modes, ω_1 is non-degenerate, ω_2 is double degenerate, and ω_3 , ω_4 , ω_5 , and ω_6 are triply degenerate. The form of these are shown in Fig. 9 in which the vibrational frequencies are labelled $\omega_1 \dots \omega_6$. ω_1 , ω_2 , and ω_3 are associated with essentially valency vibrations and ω_4 , ω_5 and ω_6 with essentially deformation vibrations. ω_1 , ω_2 , and ω_5 appear in the Raman spectrum, ω_3 and ω_4 appear in the infra-red spectrum, ω_6 does not appear in either because the transition is forbidden for both Raman and infra-red spectrum. This means that only the first five fundamental vibrational frequencies can be directly determined spectroscopically. ω_6 has to be obtained either from combination tones or from specific-heat data and it is, therefore, less accurate. With TeF_6 , ω_4 was not directly observed but was also derived from overtones. Heath and Linnett (46) feel that any value between 340 and 370 cm^{-1} would account satisfactorily for the overtones observed and that the value cannot be fixed more closely. However, more recent publications (40) and (66) assign a value of 327 cm^{-1} for this vibration.




Fig. 9 NORMAL VIBRATIONS OF AN OCTAHEDRAL XY_6 MOLECULE

Heath and Linnett (46) have examined the application of two different force fields to octahedral XY_6 molecules, and in particular TeF_6 . They found the "Orbital Valency Force Field" equations, which are based on the view that valency bonds are related to the form of the atomic orbitals, to give superior results to the "Simple Valency Force Field" equations. The following table shows a comparative agreement between observed and calculated frequencies using the OVFF and SVFF approach respectively.

Vibrational Frequencies of TeF_6 in cm^{-1}

TeF_6	ω_1	ω_2	ω_3	ω_4	ω_5	ω_6
Observed (46)	701	674	751	327	313	165
Calculated OVFF	698	669	775	351	313	388
Calculated SVFF	704	674	780	277	320	224

It is noted that the normal vibration, ω_4 , is in fairly good agreement with the calculated value; whereas the ω_6 value is not satisfactorily considered in the OVFF approach.

Heath and Linnett (46) give the following equations for TeF_6 from the OVFF approach:

$$\lambda_1 = (k_1 + 8A)/m_F \quad (\text{I.9})$$

$$\lambda_2 = (k_1 - \frac{8}{3} \frac{B}{R} + \frac{8A}{3}) m_F \quad (\text{I.10})$$

$$\begin{aligned} \lambda_3 + \lambda_4 = & f_{11}(1/m_F + 2/m_{\text{Te}}) + f_{22}(4/m_{\text{Te}} + 1/m_F) \\ & - 4 \quad 2(f_{12}/m_{\text{Te}}) \end{aligned} \quad (\text{I.11})$$

$$\lambda_3 \lambda_4 = (f_{11} f_{22} - f_{12}^2) (1/m_F^2 + 6/m_F m_{\text{Te}}) \quad (\text{I.12})$$

where,

$$f_{11} = (k_1 - \frac{2B}{R} + 4A)$$

$$f_{12} = (2 \quad 2 \quad A - 2 \quad B/R)$$

$$f_{22} = (k_B + \frac{3B}{R} + 2A)$$

$$\lambda_5 = (k_B + \frac{2B}{R} + 4A)/m_F \quad (\text{I.13})$$

$$\lambda_6 = (k_B + 2A)/m_F \quad (\text{I.14})$$

where,

$$\lambda_1 = 4\pi^2 c^2 \omega_1^2 \quad \omega_1 = \text{frequency in cm}^{-1}$$

m_F = mass of fluorine atom (amu)

m_{Te} = mass of tellurium atom (amu)

k_1 = the bond stretching constant of the
Te-F bond

k_B = the angle bending constant of the
F-Te-F bond

-B and 2A are the first and second differentials of the potential energy function representing repulsion between non-bonded atoms.

The constants have been calculated by Heath and Linnett to be:

$$k_1 = 4.85 \times 10^5 \text{ dyne/cm}$$

$$k_B = 0.78 \times 10^5 \text{ dyne/cm}$$

$$A = 0.075 \times 10^5 \text{ dyne/cm}$$

$$\frac{B}{R} = 0.0115 \times 10^5 \text{ dyne/cm}$$

The values used for the fluorine and tellurium masses are those given by Semat (94).

The first part of the paper is devoted to a discussion of the general principles of the theory of the structure of the atom. It is shown that the structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum. The structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum.

The second part of the paper is devoted to a discussion of the general principles of the theory of the structure of the atom. It is shown that the structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum. The structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum.

The third part of the paper is devoted to a discussion of the general principles of the theory of the structure of the atom. It is shown that the structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum. The structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum.

The fourth part of the paper is devoted to a discussion of the general principles of the theory of the structure of the atom. It is shown that the structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum. The structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum.

It is immediately seen from the equations that λ_1 , λ_2 , λ_5 , and λ_6 are independent of the mass of tellurium and, therefore, remain the same for isotopic substitution of tellurium. For this reason, the poor agreement of the calculated value of ω_6 with the observed value can be overlooked in calculating isotopic shifts. Using equations I.11 and I.12, $\lambda_3 + \lambda_4$ and $\lambda_3\lambda_4$ are obtained. Then, λ_3 and λ_4 and subsequently ω_3 and ω_4 can be obtained by using the equation:

$$\lambda^2 - (\lambda_3 + \lambda_4) \lambda + \lambda_3\lambda_4 = 0 \quad (\text{I.15})$$

whose roots are given by

$$2\lambda = \lambda_3 + \lambda_4 \pm \sqrt{(\lambda_3 + \lambda_4)^2 - 4\lambda_3\lambda_4} \quad (\text{I.16})$$

An IBM 7040 computer program was set up to calculate ω_3 and ω_4 for all isotopic masses of tellurium using the foregoing considerations. (See pages A63, 65, Appendix II).

The values of ω_3 and ω_4 were found to be the following:

Isotope	ω_3 cm^{-1}	ω_4 cm^{-1}
$\text{Te}^{120}\text{F}_6$	775.42	356.47
$\text{Te}^{122}\text{F}_6$	773.78	355.79
$\text{Te}^{123}\text{F}_6$	772.98	355.46
$\text{Te}^{124}\text{F}_6$	772.19	355.13
$\text{Te}^{125}\text{F}_6$	771.42	354.80
$\text{Te}^{126}\text{F}_6$	770.66	354.48
$\text{Te}^{128}\text{F}_6$	769.17	353.85
$\text{Te}^{130}\text{F}_6$	767.72	353.22

The computed values for $\text{Te}^{130}\text{F}_6$ are now compared with the observed values as follows:

$\text{Te}^{130}\text{F}_6$	ω_3 cm^{-1}	ω_4 cm^{-1}
Observed	752	327
Computed	767.72	353.22

Correction factors $\frac{752}{767.72}$ and $\frac{327}{353.22}$ are used to align the calculated vibrational frequencies for the isotopic mass with the observed vibrational frequencies for $\text{Te}^{130}\text{F}_6$.

Time	Temp	Remarks
10:00	65.0	Clear
11:00	68.0	Clear
12:00	70.0	Clear
13:00	72.0	Clear
14:00	74.0	Clear
15:00	76.0	Clear
16:00	78.0	Clear
17:00	80.0	Clear

Observer: J. H. Smith, Station: Washington, D.C.

Remarks: Clear, calm, no wind.

Time	Temp	Remarks
18:00	82.0	Clear
19:00	84.0	Clear
20:00	86.0	Clear
21:00	88.0	Clear

Time: 21:00, Temp: 88.0, Remarks: Clear

Station: Washington, D.C., Date: 10/10/1918

Observer: J. H. Smith, Station: Washington, D.C.










Then in terms of the corrected frequencies, we have :

	ω_1 cm ⁻¹	ω_2 cm ⁻¹	ω_3 cm ⁻¹	ω_4 cm ⁻¹	ω_5 cm ⁻¹	ω_6 cm ⁻¹
Te ¹²⁰ F ₆	701	674	758.52	330.00	313	164
Te ¹²² F ₆	701	674	756.92	329.38	313	164
Te ¹²³ F ₆	701	674	756.14	329.07	313	164
Te ¹²⁴ F ₆	701	674	755.37	328.77	313	164
Te ¹²⁵ F ₆	701	674	754.61	328.47	313	164
Te ¹²⁶ F ₆	701	674	753.87	328.17	313	164
Te ¹²⁸ F ₆	701	674	752.41	327.58	313	164
Te ¹³⁰ F ₆	701	674	751.00	327.00	313	164

TeBr₄

TeBr₄ belongs to the group XY₄ having a tetrahedral structure. There are four normal modes of vibration; one totally symmetric (non-degenerate) mode (ω_1) of symmetry type A₁, one doubly degenerate mode (ω_2) of type E, and two triply degenerate modes (ω_3 and ω_4) of type F₂. These normal vibrations are shown in Fig.10.

NORMAL VIBRATIONS
of a
TETRAHEDRAL XY_4 MOLECULE



Various workers have set down force equations for this molecule. Venkateswarlu, Somasundaram, and Krishna (121) give a method of evaluation, but to quote C.W.F.T. Pistorius, (85), " . . .they seem to have made a numerical error, because they list real values while the correct results according to their method undoubtedly are imaginary." I found the same to be true.

The best results were found to be those of the OVFF approach of Heath and Linnett (45). They give the force equations for some XY_4 tetrahalides as:

$$\begin{aligned}\lambda_1 &= (k + 8A)/M_F \\ \lambda_2 &= (3k/(r^0)^2 + 2A + B/R^0)M_F \\ \lambda_3 + \lambda_4 &= (k + 8A/3 - 4B/3R^0)(1/M_F + 4/3M_{Te}) \\ &\quad + (2k/(r^0)^2 + 4A/3 + 10B/3R^0)(1/M_F + 8/3M_{Te}) \\ &\quad - (8A/3 - 4B/3R^0)(8/3M_{Te}) \\ \lambda_3\lambda_4 &= (R + 8A/3 - 4B/3R^0)(2k/(r^0)^2 + \\ &\quad 4A/3 + 10B/3R^0) - 1/2(8A/3 - 4B/3R^0)^2 \\ &\quad 1/M_F^2 + 4/M_{Te} \cdot M_F\end{aligned}$$

Miss R. Thanalakshmi (103) calculated the force constants of the OVFF equations of Heath and Linnett (45) and gives the following values:

the first of these is the fact that the system is not

in equilibrium with the environment, and the second is

that the system is not in equilibrium with the environment.

The first of these is the fact that the system is not

in equilibrium with the environment, and the second is

that the system is not in equilibrium with the environment.

The first of these is the fact that the system is not

in equilibrium with the environment, and the second is

that the system is not in equilibrium with the environment.

The first of these is the fact that the system is not

in equilibrium with the environment, and the second is

that the system is not in equilibrium with the environment.

The first of these is the fact that the system is not

in equilibrium with the environment, and the second is

that the system is not in equilibrium with the environment.

The first of these is the fact that the system is not

in equilibrium with the environment, and the second is

that the system is not in equilibrium with the environment.

The first of these is the fact that the system is not

in equilibrium with the environment, and the second is

that the system is not in equilibrium with the environment.

The first of these is the fact that the system is not

in equilibrium with the environment, and the second is

that the system is not in equilibrium with the environment.

The first of these is the fact that the system is not

in equilibrium with the environment, and the second is

that the system is not in equilibrium with the environment.

$$k = 1.4500 \times 10^5 \text{ dyne/cm.}$$

$$k/(r^0)^2 = 0.0192 \times 10^5 \text{ dyne/cm.}$$

$$B/R^0 = 0.0040 \times 10^5 \text{ dyne/cm.}$$

$$A = 0.0300 \times 10^5 \text{ dyne/cm.}$$

It is noted that $\lambda_1 + \lambda_2$ are independent of the mass of tellurium and therefore remain the same for isotopic substitution of tellurium. The method of determining the roots and thus the frequencies ω_3 and ω_4 is exactly the same as was done for TeF_6 . See equations I.15 and I.16.

Again an IBM 7040 computer program was set up (Pages A63 and 65) and results are given as follows:

Isotope	ω_3 cm^{-1}	ω_4 cm^{-1}
$\text{Te}^{120}\text{Br}_4$	244.92	61.49
$\text{Te}^{122}\text{Br}_4$	243.96	61.36
$\text{Te}^{123}\text{Br}_4$	243.50	61.30
$\text{Te}^{124}\text{Br}_4$	243.04	61.23
$\text{Te}^{125}\text{Br}_4$	242.58	61.17
$\text{Te}^{126}\text{Br}_4$	242.14	61.11
$\text{Te}^{128}\text{Br}_4$	241.26	60.99
$\text{Te}^{130}\text{Br}_4$	240.41	60.86

Again a discrepancy is found between the computed and observed frequencies for $\text{Te}^{130}\text{Br}_4$

$\text{Te}^{130}\text{Br}_4$	ω_3 cm^{-1}	ω_4 cm^{-1}
observed	209	64
calculated	240.41	60.86


Upon correction, the final normal frequencies are then:

	ω_1 cm^{-1}	ω_2 cm^{-1}	ω_3 cm^{-1}	ω_4 cm^{-1}
$\text{Te}^{120}\text{Br}_4$	190	51	212.92	64.66
$\text{Te}^{122}\text{Br}_4$	190	51	212.09	64.52
$\text{Te}^{123}\text{Br}_4$	190	51	211.68	64.46
$\text{Te}^{124}\text{Br}_4$	190	51	211.29	64.39
$\text{Te}^{125}\text{Br}_4$	190	51	210.89	64.32
$\text{Te}^{126}\text{Br}_4$	190	51	210.50	64.26
$\text{Te}^{128}\text{Br}_4$	190	51	209.74	64.13
$\text{Te}^{130}\text{Br}_4$	190	51	209.00	64.00

SCTe

This molecule is of the triatomic linear type having a symmetric stretching mode ω_1 , a bending mode ω_2 , and an asymmetric stretching mode ω_3 . Of these, the

asymmetric stretching mode (ω_2) is doubly degenerate, and ω_1 and ω_3 non-degenerate (49). The forms of the normal vibrations would look like the following:



Wentink (120) gives the following force equations for this molecule:

$$\lambda_1 + \lambda_3 = K_1(1/M_S + 1/M_C) + K_2(1/M_C + 1/M_{Te}) - 2K_3/M_C$$

$$\lambda_1 \lambda_3 = \frac{M_S + M_C + M_{Te}}{M_S M_C M_{Te}} (K_1 K_2 - K_3^2)$$

$$\lambda_2 = (1/r_1^2 r_2^2) \cdot (r_1^2/M_{Te} + r_2^2/M_S + r_{12}^2/M_C) K_\alpha$$

where

K_1 = stretching constant for the S-C bond

K_2 = stretching constant for the C-Te bond

K_3 = stretching interaction constant

K_α = bending constant

r_1 = S-C internuclear distance

r_2 = C-Te internuclear distance

r_{12} = S-Te internuclear distance

The first part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation $f(x) = \int_0^x f(t) dt$. It is shown that $f(x)$ is a constant function, and its value is determined by the initial condition $f(0) = 1$.

$$\begin{aligned} f(x) &= \int_0^x f(t) dt \\ f(x) &= \int_0^x 1 dt \\ f(x) &= x \end{aligned}$$

The second part of the paper is devoted to the study of the properties of the function $g(x)$ defined by the equation $g(x) = \int_0^x g(t) dt$. It is shown that $g(x)$ is a constant function, and its value is determined by the initial condition $g(0) = 1$.

$$\begin{aligned} g(x) &= \int_0^x g(t) dt \\ g(x) &= \int_0^x 1 dt \\ g(x) &= x \end{aligned}$$

The third part of the paper is devoted to the study of the properties of the function $h(x)$ defined by the equation $h(x) = \int_0^x h(t) dt$. It is shown that $h(x)$ is a constant function, and its value is determined by the initial condition $h(0) = 1$.

$$\begin{aligned} h(x) &= \int_0^x h(t) dt \\ h(x) &= \int_0^x 1 dt \\ h(x) &= x \end{aligned}$$

The fourth part of the paper is devoted to the study of the properties of the function $i(x)$ defined by the equation $i(x) = \int_0^x i(t) dt$. It is shown that $i(x)$ is a constant function, and its value is determined by the initial condition $i(0) = 1$.

He gives the values for these constants as follows:

$$K_1 = 7.04 \times 10^5 \text{ dyne/cm.}$$

$$K_2 = 4.58 \times 10^5 \text{ dyne/cm.}$$

$$K_3 = 0.30 \times 10^5 \text{ dyne/cm.}$$

$$K_\alpha = 0.52 \times 10^{11} \text{ dyne/cm-rad}$$

$$r_1 = 1.557 \text{ Å}$$

$$r_2 = 1.904 \text{ Å}$$

$$r_{12} = 3.461 \text{ Å}$$

As noted, each λ is dependent on the mass of tellurium and therefore, each is affected by isotopic substitution. Knowing $\lambda_1 + \lambda_3$ and $\lambda_1 \lambda_3$ one uses equations I.15 and I.16 as before for obtaining λ_1 and λ_3 and subsequently ω_1 and ω_3 . λ_2 is calculated directly and from this ω_2 is obtained.

Using the computer program (pages A63, 64) the values of ω_1 , ω_2 , and ω_3 were found to be as follows:

	ω_1 cm^{-1}	ω_2 cm^{-1}	ω_3 cm^{-1}
SCTe ¹²⁰	1347.64	340.05	426.35
SCTe ¹²²	1347.54	340.00	425.45
SCTe ¹²³	1347.48	339.98	425.00
SCTe ¹²⁴	1347.43	339.95	424.57
SCTe ¹²⁵	1347.38	339.93	424.14
SCTe ¹²⁶	1347.33	339.91	423.72
SCTe ¹²⁸	1347.23	339.86	422.89
SCTe ¹³⁰	1347.14	337.00	422.08

Again a discrepancy is found between the calculated and observed frequencies for SCTe¹³⁰.

SCTe ¹³⁰	ω_1 cm^{-1}	ω_2 cm^{-1}	ω_3 cm^{-1}
Observed	1347.00	337.00	423.00
Calculated	1347.14	337.00	422.08

When this has been corrected for, the final normal frequencies are then:

Table 1			
Year	1990	1995	2000
1990	100	100	100
1991	105	105	105
1992	110	110	110
1993	115	115	115
1994	120	120	120
1995	125	125	125
1996	130	130	130
1997	135	135	135
1998	140	140	140
1999	145	145	145
2000	150	150	150

Table 1. Summary of data for the first experiment. The data are presented in the form of a table with four columns: Year, 1990, 1995, and 2000. The rows represent the years from 1990 to 2000, with the first row being the header row. The values in the table are 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, and 150, respectively.


Table 2			
Year	1990	1995	2000
1990	100	100	100
1991	105	105	105
1992	110	110	110
1993	115	115	115
1994	120	120	120
1995	125	125	125
1996	130	130	130
1997	135	135	135
1998	140	140	140
1999	145	145	145
2000	150	150	150

Table 2. Summary of data for the second experiment. The data are presented in the form of a table with four columns: Year, 1990, 1995, and 2000. The rows represent the years from 1990 to 2000, with the first row being the header row. The values in the table are 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, and 150, respectively.

	ω_1 cm ⁻¹	ω_2 cm ⁻¹	ω_3 cm ⁻¹
SCTe ¹²⁰	1347.51	340.50	427.28
SCTe ¹²⁴	1347.40	340.00	426.37
SCTe ¹²³	1347.35	339.98	425.93
SCTe ¹²⁴	1347.30	339.95	425.49
SCTe ¹²⁵	1347.24	339.93	425.06
SCTe ¹²⁶	1347.19	339.91	424.64
SCTe ¹²⁸	1347.10	339.86	423.81
SCTe ¹³⁰	1347.00	337.00	423.00

H₂Te

This molecule is of non-linear XY₂ structure having the form of normal vibrations shown below. Each vibration is non-degenerate.



Due to the chemical instability of this gas (93) high resolution spectroscopy has been limited and the vibrational frequencies given are peak values for bands of unresolved spectral lines. Some overlapping of bands is also reported.

Triatomic non-linear molecules have been studied by Glockler and Tung (41) with the assumption of general force fields which produce three equations with four force constants for isosceles non-linear triatomic molecules as follows:

$$\lambda_1 + \lambda_2 = L_{11} + L_{12} + L_{33}$$

$$\lambda_1 \lambda_2 = L_{33} (L_{11} + L_{12}) - 2L_{13}L_{31} \quad (\text{I.17})$$

$$\lambda_3 = L_{11} - L_{12}$$

where

$$-L_{11} = \frac{C_1}{\mu} + \frac{C_3}{M} \sin \alpha - \frac{C_4}{M} \cos \alpha$$

$$L_{12} = \frac{C_1}{M} \cos \alpha - \frac{C_3}{M} \sin \alpha + \frac{C_4}{\mu}$$

$$-L_{13} = \frac{C_1}{M} \sin \alpha - 2C_3 \left[\frac{1}{\mu} - \frac{\cos \alpha}{M} \right] + \frac{C_4}{M} \sin \alpha$$

$$-L_{31} = \frac{C_2}{M} \sin \alpha - C_3 \left[\frac{1}{\mu} + \frac{\cos \alpha}{M} \right]$$

$$L_{33} = 2C_2 \left[\frac{1}{\mu} - \frac{\cos \alpha}{M} \right] - \frac{2C_3}{M} \sin \alpha$$


where

M = mass of central atom

m = mass of terminal atoms

$$\frac{1}{\mu} = \frac{1}{M} + \frac{1}{m}$$

α is the apex angle as shown



The observed angle α for the H-Te bond is close to 90° (101) (Rossmann and Straley in 1956 gave it as $89^\circ 30'$ (93)). The two observed symmetric frequencies are 2047 cm^{-1} and 2133 cm^{-1} and the anti-symmetric frequency observed is 860.79 cm^{-1} (50, 102). Since α is so close to 90° , the equations are simplified to

$$-L_{11} = \frac{-C_1}{\mu} + \frac{C_3}{M}$$

$$L_{12} = \frac{-C_3}{M} + \frac{C_4}{\mu}$$

$$-L_{13} = \frac{C_1}{M} - \frac{2C_3}{\mu} + \frac{C_4}{M}$$

$$-L_{31} = \frac{C_2}{M} - \frac{C_3}{\mu}$$

$$L_{33} = \frac{2C_2}{\mu} - \frac{2C_3}{M}$$

giving

$$\lambda_1 + \lambda_2 = \frac{1}{\mu} [2c_2 + c_4 + c_1] - \frac{4c_3}{M}$$

$$\lambda_1 \lambda_2 = [2c_2(c_1 + c_4) - 4c_3^2] \left[\frac{1}{\mu^2} - \frac{1}{M^2} \right] \dots \quad (\text{I.18})$$

$$\lambda_3 = (c_1 - c_4) \left(\frac{1}{\mu} \right)$$

It is seen from I.18 that the isotopic frequencies for ω_3 are related by

$$\frac{\lambda'_3}{\lambda_3} = \frac{\mu}{\mu'}, \quad \text{or} \quad \frac{\omega'_3}{\omega_3} = \sqrt{\frac{\mu}{\mu'}}$$

and are thus independent of the force constants.

Also,

$$\frac{\lambda'_1 \lambda'_2}{\lambda_1 \lambda_2} = \frac{\left[\frac{1}{\mu'^2} - \frac{1}{M'^2} \right]}{\left[\frac{1}{\mu^2} - \frac{1}{M^2} \right]}$$

which is also independent of the force constants. Thus only the calculation of $\lambda_1 + \lambda_2$ depends on the values of the force constants and being of much smaller magnitude than the product $\lambda_1 \lambda_2$, any error in the values chosen for the force constants is very nearly negligible as affecting isotopic vibrational frequency computations.




FIG. II

It is seen by the set of equations (I.18) that only three equations are available for determining four unknown force constants. Thus it is impossible to determine the force constants uniquely. However, if C_4 is assumed known and the other constants calculated in terms of C_4 , there exists only a limited range of values of C_4 which give a 'non-imaginary' solution. The relationship between the four constants is plotted in Fig. 11. Thus the constants have a limited range of possible values, namely, $C_4 = 0.61 \pm 0.50$, $C_3 = 0 \pm 0.50$, $C_2 = 0.72 \pm 0.50$, and $C_1 = 2.05 \pm 0.50$ ($\times 10^5$ dyne/cm.). Therefore, if a unique set of force constants is to be found, one of the constants must be determined by some other means. ω_3 is the stretching frequency of the hydrogen-tellurium bond. In the polyatomic molecule its value is affected by interaction with the other forces, but one can get an approximate value for the H-Te stretching constant C_1 by considering a diatomic H-Te molecule where the interaction forces are absent. Lippincott and Dayhoff (67) give the value of ω_3 for TeH as 2250 cm^{-1} . Using the relation that

$$\omega_e = \frac{1}{2\pi c} \sqrt{\frac{k}{\mu}}$$

where c = velocity of light

μ = reduced mass

k = bond constant

k is deduced to be 2.66×10^5 dyne/cm. for the diatomic H-Te bond. This would thus delimit the values of the constants to those values appearing to the left in Fig. 10 where C_1 has the maximum value of 2.55×10^5 dyne/cm.

Smith and Linnett (99) have calculated the constants for the other members of the group VIA elements using similar considerations along with the relation:

$$[\log C_1 = -6 \log r + a]$$

from excited state spectroscopic data to deduce C_1 and therefore deduce C_2, C_3 and C_4 . They give the following set of force constants: ($\times 10^5$ dynes/cm)

	C_1	C_2	C_3	C_4
H ₂ O	8.36	0.75	0.20	-0.10
H ₂ S	4.4 $\pm$ 0.2	0.55 $\pm$ 0.05	0.2 $\pm$ 0.2	-0.15 $\pm$ 0.5
H ₂ Se	3.4 $\pm$ 0.1	0.38 $\pm$ 0.06	0.15 $\pm$ 0.03	-0.15 $\pm$ 0.15

It is noted that the value of the constants decreases with increasing order in the periodic table. Thus, this further delimits the values of the constants to $C_3 < 0.18$ which in turn limits $C_2 < 0.25$, $C_1 > 2.52$, $C_4 > -0.14$. It is noted also that C_3 is positive. In terms of this value we can then quite accurately assign the following values to the H_2Te force constants.

$$\begin{aligned} C_1 &= 2.54 \pm 0.01 \times 10^5 \text{ dyne/cm} \\ C_2 &= 0.23 \pm 0.01 \times 10^5 \text{ dyne/cm} \\ C_3 &= 0.13 \pm 0.05 \times 10^5 \text{ dyne/cm} \\ C_4 &= -0.12 \pm 0.01 \times 10^5 \text{ dyne/cm} \end{aligned}$$

Having thus determined the force constants, the isotopic frequencies are calculated and corrected to line up with the observed frequencies for H_2Te^{130} . These are as follows below:

	ω_1 cm ⁻¹	ω_2 cm ⁻¹	ω_3 cm ⁻¹
H_2Te^{120}	2047.49	861.13	2133.68
H_2Te^{122}	2047.39	861.39	2133.54
H_2Te^{123}	2047.33	861.02	2133.47
H_2Te^{124}	2047.28	860.99	2133.40
H_2Te^{125}	2047.24	860.95	2133.33
H_2Te^{126}	2047.19	860.92	2133.26
H_2Te^{128}	2047.09	860.85	2133.13
H_2Te^{130}	2047.00	860.79	2133.00

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO
 5408 S. UNIVERSITY AVE.
 CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO
 5408 S. UNIVERSITY AVE.
 CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO
 5408 S. UNIVERSITY AVE.
 CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO
 5408 S. UNIVERSITY AVE.
 CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO
 5408 S. UNIVERSITY AVE.
 CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO
 5408 S. UNIVERSITY AVE.
 CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO
 5408 S. UNIVERSITY AVE.
 CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO
 5408 S. UNIVERSITY AVE.
 CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO
 5408 S. UNIVERSITY AVE.
 CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO
 5408 S. UNIVERSITY AVE.
 CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO
 5408 S. UNIVERSITY AVE.
 CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO
 5408 S. UNIVERSITY AVE.
 CHICAGO, ILL. 60637

Tellurite ion (TeO_3^-)

The fundamental vibrational frequencies have been reported by Siebert (95) to be as follows (cm^{-1})

ω_1	ω_2	ω_3	ω_4
758	364	703	326

The tellurite ion has pyramidal XY_3 structure (see fig. 12) and a force constant analysis must take into account the apex angle of the pyramid. No experimental report of the apex angle for this ion has been reported in the literature. Therefore, the expression from Page 163 of Herzberg (49)

$$\cos^2 \beta = \frac{1}{4 \frac{\nu_3^2 \nu_4^2}{\nu_1^2 \nu_2^2} + \frac{3M_Y - M_X}{3M_Y - M_X}}$$

where $\sin \beta = \frac{2}{\sqrt{3}} \sin \frac{\alpha}{2}$

and α = apex angle

ν_i = i^{th} vibrational frequency

M_Y = mass of oxygen

M_X = mass of tellurium

was used to calculate the angle. It is noted from the expression that the angle is dependent on the tellurium mass. Therefore, α was calculated for each isotope but the computations showed a range in alpha from 101.16° in $\text{Te}^{130}\text{O}_3$ to 101.15° in Te^{122} so one could use a value of 101.15° and be well within the limits of experimental error. As to the accuracy of such a calculation, the observed angle for SeO_3 has been determined to be 100.6° (128) whereas the calculated value for $^{80}\text{SeO}_3$ was 99.7° . Assuming the mean observed angle to apply to $^{80}\text{SeO}_3$, this represents an error of less than 0.8%.

The Wilson F-G Matrix method has been used to determine the force constants. Following Nakomoto (82) the F and G matrix elements are:

F matrix elements:

For A_1 vibrations

$$\begin{aligned} F_{11} &= f_r + 2f_{rr} \\ F_{12} &= r(2f_{r\alpha} + f_{r\alpha'}) \\ F_{22} &= r^2(f_\alpha + 2f_{\alpha\alpha}) \end{aligned}$$

For E vibrations

$$\begin{aligned} F_{11} &= f_r - f_{rr} \\ F_{12} &= r(-f_{r\alpha} + f_{r\alpha'}) \\ F_{22} &= r^2(f_\alpha - f_{\alpha\alpha}) \end{aligned}$$




Fig. 12 NORMAL VIBRATIONS OF A PYRAMIDAL XY_3 MOLECULE
Taken from Nakamoto (82)

G Matrix elements:

For A_1 vibrations

$$G_{11} = \mu_y + (1 + 2 \cos \alpha) \mu_x$$

$$G_{12} = - \frac{2}{r} \frac{(1+2 \cos \alpha)(1-\cos \alpha)}{\sin \alpha} \mu_x$$

$$G_{22} = \frac{2}{r^2} \left[\frac{1+2 \cos \alpha}{1+\cos \alpha} \right] \left[\mu_y + 2\mu_x(1-\cos \alpha) \right]$$

For E_1 vibrations

$$G_{11} = \mu_y + \mu_x(1 - \cos \alpha)$$

$$G_{12} = - \frac{1}{r} \frac{(1-\cos \alpha)^2}{\sin \alpha} \mu_x$$

$$G_{22} = \frac{1}{r^2(1+\cos \alpha)} [(2+\cos \alpha)\mu_y + (1-\cos \alpha)^2 \mu_x]$$

where

$$r = \text{Te-O bond distance}$$

$$\mu_y = \frac{1}{M_{\text{Oxygen}}}$$

$$\mu_x = \frac{1}{M_{\text{Te}}}$$

The A_1 vibrations consist of the two non-degenerate vibrations ω_1 and ω_2 and the E vibrations consist of the doubly degenerate ω_3 and ω_4 vibrations.

The secular equation for both A_1 and E vibrations is

$$\lambda^2 - \lambda(F_{11}G_{11} + 2F_{12}G_{12} + F_{22}G_{22}) + (F_{11}F_{22} - F_{12}^2)(G_{11}G_{22} - G_{12}^2) = 0$$

There are six force constants but only four distinguishable frequencies so in performing the analysis it has been assumed that $f_{r\alpha}$ and $f_{r\alpha'}$ may be neglected leaving f_r , f_{rr} , f_α , and $f_{\alpha\alpha}$ to be determined. It is to be noted also, that because F_{ij} is always multiplied by G_{ij} , the "r's" cancel, so for this particular molecular structure, the Te-O bond distance need not be known if the apex angle is known. Using the frequency data and the IBM 7040 program (page A58) the following set of force constants was selected:

$$\begin{aligned} f_r &= 4.377 \times 10^5 \text{ dyne/cm.} \\ f_\alpha &= 0.493 \times 10^5 \text{ dyne/cm.} \\ f_{rr} &= 0.321 \times 10^5 \text{ dyne/cm.} \\ f_{\alpha\alpha} &= 0.078 \times 10^5 \text{ dyne/cm.} \end{aligned}$$

THE HISTORY OF THE

REIGN OF

CHARLES THE FIRST

BY
JOHN BURNET
OF
GLASGOW
IN
SCOTLAND
BY
JOHN BURNET
OF
GLASGOW
IN
SCOTLAND
BY
JOHN BURNET
OF
GLASGOW
IN
SCOTLAND

THE HISTORY OF THE
REIGN OF
CHARLES THE FIRST
BY
JOHN BURNET
OF
GLASGOW
IN
SCOTLAND

These were then used to calculate the isotopic frequencies given below:

	ω_1 cm^{-1}	ω_2 cm^{-1}	ω_3 cm^{-1}	ω_4 cm^{-1}
Te ¹²⁰	760.56	366.97	706.89	326.79
Te ¹²²	760.01	366.35	706.07	326.62
Te ¹²³	759.74	366.04	705.66	326.54
Te ¹²⁴	759.48	365.74	705.26	326.46
Te ¹²⁵	759.22	365.44	704.87	326.38
Te ¹²⁶	759.97	365.14	704.49	326.30
Te ¹²⁸	758.40	364.56	703.73	326.15
Te ¹³⁰	758.00	364.00	703.10	326.00

Date		Description		Amount
1890	Jan 1	Balance		100.00
	Feb 1	Interest		5.00
	Mar 1	Interest		5.00
	Apr 1	Interest		5.00
	May 1	Interest		5.00
	Jun 1	Interest		5.00
	Jul 1	Interest		5.00
	Aug 1	Interest		5.00
	Sep 1	Interest		5.00
	Oct 1	Interest		5.00
	Nov 1	Interest		5.00
	Dec 1	Interest		5.00
	Total			100.00

APPENDIX II

IBM 7040 computer programs for
calculation of isotopic vibrational
frequencies and partition function
ratios.



A. Calculation of isotopic vibrational frequencies and partition function ratios for diatomic species

In this program, ω_{heavy} and ω_{light} are computed for diatomic species for which values of ω_e and $\omega_e x_e$ are available. The partition function ratio for temperatures $0^\circ - 100^\circ$ in steps of 10° and from $100^\circ - 1000^\circ\text{C.}$ in steps of 100° is then computed according to equation 15 on page 48.

The following symbols were employed for data which must be supplied:

$X130, X122$ = masses of the two Y-atom isotopes for which isotopic data is desired (atomic weight units)

$WE130(I)$ = known value of ω_e for the I^{th} diatomic molecule applying to the most abundant Y isotope, in this case Te^{130} .

$WEX130(I)$ = known value of $\omega_e x_e$ for the I^{th} molecule and most abundant isotope.

$X1(I)$ = mass of the I^{th} X atom in XY diatomic molecules, eg., Ge^{74} (awu).

In this work, the ratio of $\text{Te}^{130}/\text{Te}^{122}$ was needed. Te^{130} is the most abundant isotope so computation of ω_e and $\omega_e x_e$ for the heavy isotope was unnecessary. Should the heavy isotope not be the most abundant, then statements must be included computing ω_e and $\omega_e x_e$ for the heavy isotope.

\$IRSYS

\$JOB 735003 PARTITION FUNCTIONS DIATOMIC

\$TIME 003

\$IBJOB TF NODECK

\$IBFTC DIATOM NOLIST

DIMENSION WE130(7), WEX130(7), RM130(7), RM122(7), X1(7), P(7),
1SQRP(7), WE122(7), WEX122(7), W130(7), W122(7), T(20), UU(20),
2U130(7,20), U122(7,20), A(7,20), B(7,20), PR(7,20)

X130=129.94853

X122=121.94193

WE130(1)=323.4

WE130(2)=211.8

WE130(3)=250.5

WE130(4)=796.0

WE130(5)=480.4

WE130(6)=251.5

WE130(7)=2250.0

WEX130(1)=1.00

WEX130(2)=0.12

WEX130(3)=0.50

WEX130(4)=3.50

WEX130(5)=1.16

WEX130(6)=0.55

WEX130(7)=41.3

X1(1)=73.9426

X1(2)=208.04754

X1(3)=119.94059

X1(4)=16.

X1(5)=27.985825

X1(6)=129.94853

X1(7)=1.008

DO 20 I=1,7

RM130(I)=X1(I)*X130/(X1(I)+X130)

RM122(I)=X1(I)*X122/(X1(I)+X122)

SQRP(I)=RM130(I)/RM122(I)

P(I)=SQRP(I)**0.5

WE122(I)=P(I)*WE130(I)

WEX122(I)=SQRP(I)*WEX130(I)

W130(I)=WE130(I)-2.*WEX130(I)

20 W122(I)=WE122(I)-2.*WEX122(I)

DO 28 I=2,11

T(1)=0.

28 T(I)=T(I-1)+10.

DO 29 I=12,20

29 T(I)=T(I-1)+100.

E=2.71828

V=1.438828

THE JOURNAL OF THE

1900

1901

1902

1903

1904

1905

1906

1907

1908

1909

1910

1911

1912

1913

1914

1915

1916

1917

1918

1919

1920

1921

1922

1923

1924

1925

1926

1927

1928

1929

1930

1931

1932

1933

1934

1935


```

DO 21 I=1,7
DO 21 J=1,20
UU(J)=V/(T(J)+273.)
U130(I,J)=UU(J)*W130(I)
U122(I,J)=UU(J)*W122(I)
A(I,J)=-U130(I,J)/2.
B(I,J)=-U122(I,J)/2.
21 PR(I,J)=(U130(I,J)*(E**A(I,J))*(1.-(F**(2.*B(I,J))))
1/(U122(I,J)*(E**B(I,J))*(1.-(F**(2.*A(I,J))))))
WRITE (6,4)
WRITE (6,3) (SQRP(I), I=1,7)
WRITE (6,5)
WRITE (6,3) (P(I), I=1,7)
WRITE (6,6)
WRITE (6,3) (WF122(I), I=1,7)
WRITE (6,7)
WRITE (6,3) (WEX122(I), I=1,7)
WRITE (6,8)
WRITE (6,3) (W130(I), I=1,7)
WRITE (6,9)
WRITE (6,3) (W122(I), I=1,7)
WRITE (6,11)
DO 10 J=1,20
10 WRITE (6,12) (PR(I,J), I=1,7)
3 FORMAT (2X,F15.8)
4 FORMAT (18H THIS IS U130/U122)
5 FORMAT (10H THIS IS P)
6 FORMAT (14H THIS IS WF122)
7 FORMAT (16H THIS IS WFXE122)
8 FORMAT (13H THIS IS W130)
9 FORMAT (13H THIS IS W122)
11 FORMAT (40H THESE ARE THE PARTITION FUNCTION RATIOS)
12 FORMAT (7(2X,F15.8))
CALL EXIT
END
$ENTRY DIATOM

```


B. Computation of isotopic vibrational frequencies for
XY₆ molecules

This program was set up to compute isotopic vibrational frequencies according to the OVFF equations of Heath and Linnett (46). The masses of the (I) isotopes (awu) for which vibrational data is desired are used in the program as data cards. The following symbols apply to data which must be supplied:

AK = force constant k_1 (dynes/cm)

BK = force constant k_β

A = force constant A

C = force constant B/R

YM = mass of the Y atom (awu)


```

$IRSYS
$JOB          735003  VIBRATIONAL WAVELENGTHS XY6
$IRJOB TFF6    NODECK
$IRBTC OCTA    NOLIST
      DIMENSION XM(8),H(8),P(8),Q1(8),Q2(8),
1 W1(8),W2(8),CW1(8),CW2(8)
      AK=4.85E+05
      BK=0.78E+05
      A=0.075E+05
      C=0.0115E+05
      YM=19.004444
      F11=(AK-2.*C+4.*A)
      F12=(2.8284*A-1.4142*C)
      F22=(BK+3.*C+2.*A)
      READ (5,1) N
      DO 22 I=1,N
      READ (5,2) XM(I)
      H(I)=(F11)*(1./YM+2./XM(I))+F22*(4./XM(I)+1./YM)
1-5.6568*F12/XM(I)
      P(I)=(F11*F22-F12**2)*(1./YM**2+6./(YM*XM(I)))
      Q1(I)=.5*(H(I)+SQRT (H(I)**2-4.*P(I)))
      Q2(I)=.5*(H(I)-SQRT (H(I)**2-4.*P(I)))
      W1(I)=SQRT (Q1(I)/.058894)
22 W2(I)=SQRT (Q2(I)/.058894)
      DO 30 I=1,N
      CW1(I)=751.*W1(I)/W1(8)
30 CW2(I)=327.*W2(I)/W2(8)
      WRITE (6,3)
      WRITE (6,11) (H(I),I=1,N)
      WRITE (6,4)
      WRITE (6,11) (P(I),I=1,N)
      WRITE (6,5)
      WRITE (6,11) (Q1(I),I=1,N)
      WRITE (6,6)
      WRITE (6,11) (Q2(I),I=1,N)
      WRITE (6,7)
      WRITE (6,11) (W1(I),I=1,N)
      WRITE (6,8)
      WRITE (6,11) (W2(I),I=1,N)
      WRITE (6,9)
      WRITE (6,11) (CW1(I),I=1,N)
      WRITE (6,10)
      WRITE (6,11) (CW2(I),I=1,N)
1 FORMAT (I4)
2 FORMAT (2X,F10.5)
3 FORMAT (30H THESE ARE THE WAVELENGTH SUMS)
4 FORMAT (34H THESE ARE THE WAVELENGTH PRODUCTS)
5 FORMAT (24H THESE ARE THE SUM ROOTS)
6 FORMAT (31H THESE ARE THE DIFFERENCE ROOTS)

```



```
7 FORMAT (31H THIS IS ONE SET OF WAVLENGTHS)
8 FORMAT (37H THIS IS THE OTHER SET OF WAVLENGTHS)
9 FORMAT (28H THESE ARE THE CORRECTED W1S)
10 FORMAT (28H THESE ARE THE CORRECTED W2S)
11 FORMAT (2X,E14.8)
    CALL EXIT
    END
```

\$ENTRY OCTA

```
8
119.94288
121.94193
122.94368
123.94278
124.94460
125.94420
127.94649
129.94853
```


C. Computation of isotopic vibrational frequencies for
 XY_4 molecules

This program is a solution of the OVFF equations of Heath and Linnett (45). In the program, the mass of the (I) X isotopes (awu) for which frequency data is desired are put in as data cards to be read and,

$$B = K$$

$$A = A$$

$$Z = 2k/(r_o)^2$$

$$YM = \text{mass of the Y atom (awu)}$$


```

$IBSYS
$JOB          735003      VIBRATIONAL WAVELENGTHS TETRAHEDRAL
$IRJOB  TEBR4      NODECK
$IRFTC  TETRA      NOLIST
      DIMENSION XM(8),H(8),P(8),Q1(8),Q2(8),
      1W1(8),W2(8),CW1(8),CW2(8)
      R=145.0F3
      A=3.0E3
      Z=0.0384E+05
      W=400.0
      YM=79.916
      READ (5,1) N
      DO 22 I=1,N
      READ (5,2) XM(I)
      H(I)=(B+2.6666667*A-1.3333333*W)*(1./YM+1.3333333/XM(I))
      S+(1./YM+2.6666667/XM(I))*(Z+1.3333333*A+3.333333*W)
      S-(2.6666667*A-1.3333333*W)*(2.6666667/XM(I))
      P(I)=(1./(YM**2)+4./(XM(I)*YM))*(B+2.6666667*A-1.3333333*W)
      S*(Z+1.3333333*A+3.3333333*W)-.5*(2.6666667*A-1.3333333*W)**2)
      Q1(I)=.5*(H(I)+SQRT (H(I)**2-4.*P(I)))
      Q2(I)=.5*(H(I)-SQRT (H(I)**2-4.*P(I)))
      W1(I)=SQRT (Q1(I)/.058894)
22  W2(I)=SQRT (Q2(I)/.058894)
      DO 30 I=1,N
      CW1(I)=209.*W1(I)/W1(8)
30  CW2(I)=64.*W2(I)/W2(8)
      WRITE (6,3)
      WRITE (6,11) (H(I),I=1,N)
      WRITE (6,4)
      WRITE (6,11) (P(I),I=1,N)
      WRITE (6,5)
      WRITE (6,11) (Q1(I),I=1,N)
      WRITE (6,6)
      WRITE (6,11) (Q2(I),I=1,N)
      WRITE (6,7)
      WRITE (6,11) (W1(I),I=1,N)
      WRITE (6,8)
      WRITE (6,11) (W2(I),I=1,N)
      WRITE (6,9)
      WRITE (6,11) (CW1(I),I=1,N)
      WRITE (6,10)
      WRITE (6,11) (CW2(I),I=1,N)
1  FORMAT (I4)
2  FORMAT (2X,F10.5)
3  FORMAT (30H THESE ARE THE WAVELENGTH SUMS)
4  FORMAT (34H THESE ARE THE WAVELENGTH PRODUCTS)
5  FORMAT (24H THESE ARE THE SUM ROOTS)
6  FORMAT (31H THESE ARE THE DIFFERENCE ROOTS)

```



```
7 FORMAT (31H THIS IS ONE SET OF WAVELENGTHS)
8 FORMAT (37H THIS IS THE OTHER SET OF WAVELENGTHS)
9 FORMAT (28H THESE ARE THE CORRECTED W1S)
10 FORMAT (28H THESE ARE THE CORRECTED W2S)
11 FORMAT (2X,F14.8)
    CALL EXIT
    END
```

```
$ENTRY          TETRA
```

```
8
```

```
119.94288
```

```
121.94193
```

```
122.94368
```

```
123.94278
```

```
124.94460
```

```
125.94420
```

```
127.94649
```

```
129.94853
```


D. Computation of vibrational frequencies for linear XYZ molecules

This program computes the isotopic vibrational frequencies using the force field equations and force constants of Wentink (138). The (I) X-isotopic masses for which vibrational frequencies are desired are put into the program and read as data cards. The symbols used are the following:

$$A = k_1 \text{ (dynes/cm)}$$

$$B = k_2$$

$$C = k_3$$

$$D = k_\alpha$$

$$E = r_1 \text{ (cm)}$$

$$F = r_2$$

$$G = r_{12}$$

$$YM = \text{mass of the Y atom (amu)}$$

The program computes k_α for the most abundant isotope (in this case the eighth). If a known value of k_α is desired to be used, instead of using $D = A^2/DM$, put $D = (\text{value of } k_\alpha)$ and remove statements for A^2 , R , and DM .


```

$IBSYS
$JOB          735003  VIBRATIONAL WAVELENGTHS TRIATOMIC LINEAR
$IBJOB TRI    NODECK
$IBFTC SCTE   NOLIST
      DIMENSION XM(8),H(8),P(8),Q1(8),Q2(8),Q3(8),
1 W1(8),W2(8),W3(8),CW1(8),CW2(8),CW3(8)
      A=704.0F3
      R=458.0F3
      C=30.0F3
      R=337.
      A2=0.058894*(R**2)
      E=1557.0E-11
      F=1904.0E-11
      G=3461.0E-11
      ZM=31.98
      YM=12.
      READ (5,1) N
      DO 32 I=1,N
      READ (5,2) XM(I)
      DM=((F**2/XM(8))+(F**2/ZM)+(G**2/YM))/((E**2)*(F**2))
      D=A2/DM
      H(I)=A*(1./ZM+1./YM)+R*(1./YM+1./XM(I))-2.*C/YM
      P(I)=(ZM+YM+XM(I))*((A*R)-C**2)/(ZM*YM*XM(I))
      Q2(I)=((F**2/XM(I))+(F**2/ZM)+(G**2/YM))*D/((E**2)*(F**2))
      Q1(I)=.5*(H(I)+SQRT(H(I)**2-4.*P(I)))
      Q3(I)=.5*(H(I)-SQRT(H(I)**2-4.*P(I)))
      W1(I)=SQRT(Q1(I)/0.058894)
      W2(I)=SQRT(Q2(I)/0.058894)
32  W3(I)=SQRT(Q3(I)/0.058894)
      DO 33 I=1,N
      CW1(I)=1347.*W1(I)/W1(8)
      CW2(I)=337.*W2(I)/W2(8)
33  CW3(I)=423.*W3(I)/W3(8)
      WRITE (6,3)
      WRITE (6,14) (H(I),I=1,N)
      WRITE (6,4)
      WRITE (6,14) (P(I),I=1,N)
      WRITE (6,5)
      WRITE (6,14) (Q2(I),I=1,N)
      WRITE (6,6)
      WRITE (6,14) (Q1(I),I=1,N)
      WRITE (6,7)
      WRITE (6,14) (Q3(I),I=1,N)
      WRITE (6,8)
      WRITE (6,14) (W1(I),I=1,N)

```



```

WRITE (6,9)
WRITE (6,14) (W2(I),I=1,N)
WRITE (6,10)
WRITE (6,14) (W3(I),I=1,N)
WRITE (6,11)
WRITE (6,14) (CW1(I),I=1,N)
WRITE (6,12)
WRITE (6,14) (CW2(I),I=1,N)
WRITE (6,13)
WRITE (6,14) (CW3(I),I=1,N)
WRITE (6,15)
WRITE (6,14) D
1 FORMAT (I4)
2 FORMAT (2X,F10.5)
3 FORMAT (30H THESE ARE THE WAVELENGTH SUMS)
4 FORMAT (34H THESE ARE THE WAVELENGTH PRODUCTS)
5 FORMAT (11H THIS IS Q2)
6 FORMAT (24H THESE ARE THE SUM ROOTS)
7 FORMAT (31H THESE ARE THE DIFFERENCE ROOTS)
8 FORMAT (11H THIS IS W1)
9 FORMAT (11H THIS IS W2)
10 FORMAT (11H THIS IS W3)
11 FORMAT (28H THESE ARE THE CORRECTED W1S)
12 FORMAT (28H THESE ARE THE CORRECTED W2S)
13 FORMAT (28H THESE ARE THE CORRECTED W3S)
14 FORMAT (2X,E15.8)
15 FORMAT (16H THIS IS K-ALPHA)
CALL EXIT
END
$ENTRY          SCTE
8
119.94288
121.94193
122.94368
123.94278
124.94460
125.94420
127.94649
129.94853

```

1	1000	1000	1000
2	1000	1000	1000
3	1000	1000	1000
4	1000	1000	1000
5	1000	1000	1000
6	1000	1000	1000
7	1000	1000	1000
8	1000	1000	1000
9	1000	1000	1000
10	1000	1000	1000
11	1000	1000	1000
12	1000	1000	1000
13	1000	1000	1000
14	1000	1000	1000
15	1000	1000	1000
16	1000	1000	1000
17	1000	1000	1000
18	1000	1000	1000
19	1000	1000	1000
20	1000	1000	1000
21	1000	1000	1000
22	1000	1000	1000
23	1000	1000	1000
24	1000	1000	1000
25	1000	1000	1000
26	1000	1000	1000
27	1000	1000	1000
28	1000	1000	1000
29	1000	1000	1000
30	1000	1000	1000
31	1000	1000	1000
32	1000	1000	1000
33	1000	1000	1000
34	1000	1000	1000
35	1000	1000	1000
36	1000	1000	1000
37	1000	1000	1000
38	1000	1000	1000
39	1000	1000	1000
40	1000	1000	1000
41	1000	1000	1000
42	1000	1000	1000
43	1000	1000	1000
44	1000	1000	1000
45	1000	1000	1000
46	1000	1000	1000
47	1000	1000	1000
48	1000	1000	1000
49	1000	1000	1000
50	1000	1000	1000
51	1000	1000	1000
52	1000	1000	1000
53	1000	1000	1000
54	1000	1000	1000
55	1000	1000	1000
56	1000	1000	1000
57	1000	1000	1000
58	1000	1000	1000
59	1000	1000	1000
60	1000	1000	1000
61	1000	1000	1000
62	1000	1000	1000
63	1000	1000	1000
64	1000	1000	1000
65	1000	1000	1000
66	1000	1000	1000
67	1000	1000	1000
68	1000	1000	1000
69	1000	1000	1000
70	1000	1000	1000
71	1000	1000	1000
72	1000	1000	1000
73	1000	1000	1000
74	1000	1000	1000
75	1000	1000	1000
76	1000	1000	1000
77	1000	1000	1000
78	1000	1000	1000
79	1000	1000	1000
80	1000	1000	1000
81	1000	1000	1000
82	1000	1000	1000
83	1000	1000	1000
84	1000	1000	1000
85	1000	1000	1000
86	1000	1000	1000
87	1000	1000	1000
88	1000	1000	1000
89	1000	1000	1000
90	1000	1000	1000
91	1000	1000	1000
92	1000	1000	1000
93	1000	1000	1000
94	1000	1000	1000
95	1000	1000	1000
96	1000	1000	1000
97	1000	1000	1000
98	1000	1000	1000
99	1000	1000	1000
100	1000	1000	1000

E. Computation of force constants for non-linear XY_2 molecules

The force equations used are those given by Glockler and Tung (41). Only three equations are available to determine four force constants. Therefore, programs were set up to determine (1) the possible range of values of the constant C_4 (other values give imaginary results), and (2) the values of the other three constants for values of the permissible range of C_4 .

Program (1) gives the lower limit to the permissible range of C_4 , and program (2) gives the upper limit.

In these programs,

TM = mass of the most abundant isotope to which the observed fundamental frequencies apply (grams).

U = the reduced mass of the X-Y atom
ie., $1/u = 1/M_x + 1/M_y$

$$A1 = \lambda_1 + \lambda_2$$

$$A2 = \lambda_1 \lambda_2$$

$$A3 = \lambda_3$$

$C4$ = force constant C_4 (dynes/cm)

The observed fundamental vibrational frequencies are put in as data cards to be read in order $\omega_1, \omega_2, \omega_3$ (cm^{-1}).

THE UNIVERSITY OF CHICAGO PRESS

CHICAGO, ILL.

THE UNIVERSITY OF CHICAGO PRESS
1207 EAST 59TH STREET
CHICAGO, ILL. 60637
U.S.A.
LONDON: ROUTLEDGE Kegan Paul
11 BEDFORD SQUARE
LONDON, W.C.1A 3EF
ENGLAND
TOKYO: Rinsen Shoten
2-2-1 Nishi-Shinjuku
SHINJUKU-KU
TOKYO 163
JAPAN
SINGAPORE: Raffles Book Company
110 NORTH BRIDGE ROAD
SINGAPORE 205
MALAYSIA: Raffles Book Company
110 NORTH BRIDGE ROAD
SINGAPORE 205
AUSTRALIA: Raffles Book Company
110 NORTH BRIDGE ROAD
SINGAPORE 205

THE UNIVERSITY OF CHICAGO PRESS
1207 EAST 59TH STREET
CHICAGO, ILL. 60637
U.S.A.
LONDON: ROUTLEDGE Kegan Paul
11 BEDFORD SQUARE
LONDON, W.C.1A 3EF
ENGLAND
TOKYO: Rinsen Shoten
2-2-1 Nishi-Shinjuku
SHINJUKU-KU
TOKYO 163
JAPAN
SINGAPORE: Raffles Book Company
110 NORTH BRIDGE ROAD
SINGAPORE 205
MALAYSIA: Raffles Book Company
110 NORTH BRIDGE ROAD
SINGAPORE 205
AUSTRALIA: Raffles Book Company
110 NORTH BRIDGE ROAD
SINGAPORE 205

THE UNIVERSITY OF CHICAGO PRESS
1207 EAST 59TH STREET
CHICAGO, ILL. 60637
U.S.A.
LONDON: ROUTLEDGE Kegan Paul
11 BEDFORD SQUARE
LONDON, W.C.1A 3EF
ENGLAND
TOKYO: Rinsen Shoten
2-2-1 Nishi-Shinjuku
SHINJUKU-KU
TOKYO 163
JAPAN
SINGAPORE: Raffles Book Company
110 NORTH BRIDGE ROAD
SINGAPORE 205
MALAYSIA: Raffles Book Company
110 NORTH BRIDGE ROAD
SINGAPORE 205
AUSTRALIA: Raffles Book Company
110 NORTH BRIDGE ROAD
SINGAPORE 205

THE UNIVERSITY OF CHICAGO PRESS
1207 EAST 59TH STREET
CHICAGO, ILL. 60637
U.S.A.
LONDON: ROUTLEDGE Kegan Paul
11 BEDFORD SQUARE
LONDON, W.C.1A 3EF
ENGLAND
TOKYO: Rinsen Shoten
2-2-1 Nishi-Shinjuku
SHINJUKU-KU
TOKYO 163
JAPAN
SINGAPORE: Raffles Book Company
110 NORTH BRIDGE ROAD
SINGAPORE 205
MALAYSIA: Raffles Book Company
110 NORTH BRIDGE ROAD
SINGAPORE 205
AUSTRALIA: Raffles Book Company
110 NORTH BRIDGE ROAD
SINGAPORE 205

```

$IBSYS
$JOB          735003          FORCE CONSTANTS TRIATOMIC (1)
$IBJOB H2TE      NODECK
$IBFTC FORCON  NOLIST
      DIMENSION WW(3)
      TM=130.*(1./(6.025*(10.**3)))
      U=1./((6.025*(10.**3))*((1./130.)+1.))
      READ (5,1)N
      DO 22 I=1,N
22  READ (5,2) WW(I)
      A1=355.306*(WW(1)**2+WW(2)**2)
      A2=355.306**2*(WW(1)**2*WW(2)**2)
      A3=355.306*(WW(3)**2)
      B=A1-A3
      D=A2/(1./(U**2)-1./(TM**2))
      E=D/2.
      P=(2.*(U**2)*A3)/TM
      R=(4.*U)/TM
      S=((U**2)*A3*B)/2.
      T=((2.*P*R)/8.)-(U*A3)+(U*B)
      V=((R**2)/8.)-2.
      X=((P**2)/8.)-S
      C4=((-T-((T**2)-(4.*X*V))**0.5)/(2.*V))
      WRITE (6,9)
      WRITE (6,7) C4
1  FORMAT (I4)
2  FORMAT (2X,F10.5)
7  FORMAT (2X,E15.8)
9  FORMAT (11H THIS IS C4)
      CALL EXIT
      END
$ENTRY          FORCON
3
2047.
860.79
2133.

```



```

$IBSYS
$JOB          735 03          FORCE CONSTANTS TRIATOMIC (2)
$IBJOB H2TE      NODECK
$IBFTC FORCON  NOLIST
      DIMENSION WW(3)
      TM=130.*(1./((6.025*(10.**3)))
      U=1./(((6.025*(10.**3))*((1./130.)+1.))
      READ (5,1)N
      DO 22 I=1,N
22  READ (5,2) WW(I)
      A1=355.306*(WW(1)**2+WW(2)**2)
      A2=355.306**2*(WW(1)**2*WW(2)**2)
      A3=355.306*(WW(3)**2)
      B=A1-A3
      D=A2/(1./(U**2)-1./(TM**2))
      E=D/2.
      P=(2.*(U**2)*A3)/TM
      R=(4.*U)/TM
      S=((U**2)*A3*R)/2.
      T=((2.*P*R)/8.)-(U*A3)+(U*B)
      V=((R**2)/8.)-2.
      X=((P**2)/8.)-S
      C4=((-T+((T**2)-(4.*X*V))**0.5)/(2.*V))+(0.01*10.**5)
      WRITE (6,9)
      WRITE (6,7) C4
1  FORMAT (I4)
2  FORMAT (2X,F10.5)
7  FORMAT (2X,E15.8)
9  FORMAT (11H THIS IS C4)
      CALL EXIT
      END
$ENTRY          FORCON
3
2047.
860.79
2133.

```


Programs (3) and (4) then give the values of the constants C_1 , C_2 , and C_3 for the permissible range of C_4 , C_4 going in steps of 0.01×10^5 from maximum value to minimum value. Two programs are necessary, since for each value of C_4 , C_3 has values

$$\frac{H \pm \sqrt{H^2 - 8G}}{4}$$

where H and G are defined as in the program.

Defining the symbols,

TM = mass of the most abundant isotope to

which the observed fundamental frequencies
apply (grams)

U, A1, A2, A3, C4 as previously defined

C1 = force constant C_1 (dynes/cm)

C32 = force constant C_3

C22 = force constant C_2

The observed fundamental frequencies are again
'read in as data cards in order.

THE UNIVERSITY OF CHICAGO

IN THE DEPARTMENT OF THE HISTORY OF ARTS AND ARCHITECTURE

A DISSERTATION SUBMITTED TO THE FACULTY OF THE DIVISION OF THE PHYSICAL SCIENCES

IN CANDIDACY FOR THE DEGREE OF DOCTOR OF PHILOSOPHY

BY

JOHN H. COOPER

1967

CHICAGO, ILLINOIS

UNIVERSITY OF CHICAGO PRESS

Copyright © 1967 by The University of Chicago Press

Printed in the United States of America

Library of Congress Catalog Card No. 67-21111

First published 1967

Reprinted 1970, 1971, 1972, 1973, 1974, 1975, 1976, 1977, 1978, 1979, 1980, 1981, 1982, 1983, 1984, 1985, 1986, 1987, 1988, 1989, 1990, 1991, 1992, 1993, 1994, 1995, 1996, 1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025

ISBN 0-226-03411-1

0-226-03411-1

0-226-03411-1

0-226-03411-1

```

$IRSYS
$JOB          735003          FORCE CONSTANTS TRIATOMIC (3)
$TIME         005
$IBJOB H2TE    NODECK
$IBFTC FORCON  NOLIST
    DIMENSION WW(3)
    TM=130.*(1./(6.025*(10.**3)))
    U=1./((6.025*(10.**3))*((1./130.)+1.))
    READ (5,1)N
    DO 22 I=1,N
22  READ (5,2) WW(I)
    A1=355.306*(WW(1)**2+WW(2)**2)
    A2=355.306**2*(WW(1)**2*WW(2)**2)
    A3=355.306*(WW(3)**2)
    B=A1-A3
    D=A2/(1./(U**2)-1./(TM**2))
    E=D/2.
    P=(2.*(U**2)*A3)/TM
    R=(4.*U)/TM
    S=((U**2)*A3*B)/2.
    T=((2.*P*R)/8.)-(U*A3)+(U*B)
    V=((P**2)/8.)-2.
    X=((P**2)/8.)-S
    C4=-0.05*10.**5
10  C1=U*A3+C4
    F=B/2.-C4/U
    G=F-((U**2)*A3*F)-2.*U*F*C4
    H=(2.*(U**2)*A3/TM+4.*U*C4/TM)
    C32=(H-((H**2)-8.*G)**0.5)/4.
    C22=U*(F+(2.*C32/TM))
    WRITE (6,9)
    WRITE (6,7) C4
    WRITE (6,4)
    WRITE (6,7) C32
    WRITE (6,6)
    WRITE (6,7) C22
    WRITE (6,8)
    WRITE (6,7) C1
    C4=C4-(0.01*10.**5)
    IF(C4.GE.-1.15*10.**5) GO TO 10
1  FORMAT (I4)
2  FORMAT (2X,F10.5)
4  FORMAT (11H THIS IS C3)
6  FORMAT (11H THIS IS C2)
7  FORMAT (2X,E15.8)
8  FORMAT (11H THIS IS C1)
9  FORMAT (11H THIS IS C4)
    CALL EXIT
    END
$ENTRY          FORCON
3
2047.
860.79
2133.

```



```

$IRSYS
$JOB          735003          FORCE CONSTANTS TRIATOMIC (4)
$TIME         005
$IBJOB H2TE     NODECK
$IBFTC FORCON  NOLIST
      DIMENSION WW(3)
      TM=130.*(1./(6.025*(10.**3)))
      U=1./(((6.025*(10.**3))*((1./130.)+1.))
      READ (5,1)N
      DO 22 I=1,N
22  READ (5,2) WW(I)
      A1=355.306*(WW(1)**2+WW(2)**2)
      A2=355.306**2*(WW(1)**2*WW(2)**2)
      A3=355.306*(WW(3)**2)
      B=A1-A3
      D=A2/(1./(U**2)-1./(TM**2))
      E=D/2.
      P=(2.*(U**2)*A3)/TM
      R=(4.*U)/TM
      S=((U**2)*A3*B)/2.
      T=((2.*P*R)/8.)-(U*A3)+(U*B)
      V=((P**2)/8.)-2.
      X=((P**2)/8.)-S
      C4=-0.05*10.**5
10  C1=U*A3+C4
      F=B/2.-C4/U
      G=E-((U**2)*A3*F)-2.*U*F*C4
      H=(2.*(U**2)*A3/TM+4.*U*C4/TM)
      C32=(H+((H**2)-8.*G)**0.5)/4.
      C22=U*(F+(2.*C32/TM))
      WRITE (6,9)
      WRITE (6,7) C4
      WRITE (6,4)
      WRITE (6,7) C32
      WRITE (6,6)
      WRITE (6,7) C22
      WRITE (6,8)
      WRITE (6,7) C1
      C4=C4-(0.01*10.**5)
      IF(C4.GE.-1.15*10.**5) GO TO 10
1  FORMAT (I4)
2  FORMAT (2X,F10.5)
4  FORMAT (11H THIS IS C3)
6  FORMAT (11H THIS IS C2)
7  FORMAT (2X,F15.8)
8  FORMAT (11H THIS IS C1)
9  FORMAT (11H THIS IS C4)
      CALL EXIT
      END
$ENTRY          FORCON
3
2047.
860.79
2133.

```


F. Computation of isotopic vibrational frequencies and partition function ratios for non-linear XY_2 molecules

Knowing the value of the four force constants $C_1 \dots C_4$, this program computes the isotopic vibrational frequencies and partition function ratios for non-linear XY_2 molecules according to the equations given by Glockler and Tung (41).

The partition function ratios are given for temperatures $0^\circ - 100^\circ\text{C}$ in steps of 10° and from $100^\circ - 1000^\circ\text{C}$ in steps of 100° . The isotopic masses (awu) are read in as data cards, and symbols for which data must be supplied are:

$W(I) = \omega(I)$ observed (cm^{-1})

CI = force constants $C_1 \dots C_4$ (dyne/cm)

YM = mass of the Y atom (awu)

DEGI = degree of degeneracy of the I^{th} vibration

$U(L \text{ etc}) = hc \cdot CWW(I,J)/kT(L)$ where $CWW(I,J)$ is the corrected I^{th} frequency of the J^{th} isotope for which the partition function ratio is desired

The partition function ratio is computed according to equation 16, page 49 and will be the ratio for those masses included in the $U(L)$ statements, eg. $CWW(I,1)$ will give the ratio of the 7^{th} isotope to the first (see page A58)

The first part of the paper is devoted to a general discussion of the problem of the origin of the universe. It is shown that the question of the origin of the universe is a question of the origin of the material world. The question of the origin of the material world is a question of the origin of the matter. The question of the origin of the matter is a question of the origin of the atoms. The question of the origin of the atoms is a question of the origin of the molecules. The question of the origin of the molecules is a question of the origin of the cells. The question of the origin of the cells is a question of the origin of the organisms. The question of the origin of the organisms is a question of the origin of the species. The question of the origin of the species is a question of the origin of the races. The question of the origin of the races is a question of the origin of the nations. The question of the origin of the nations is a question of the origin of the states. The question of the origin of the states is a question of the origin of the governments. The question of the origin of the governments is a question of the origin of the laws. The question of the origin of the laws is a question of the origin of the customs. The question of the origin of the customs is a question of the origin of the habits. The question of the origin of the habits is a question of the origin of the character. The question of the origin of the character is a question of the origin of the soul. The question of the origin of the soul is a question of the origin of the spirit. The question of the origin of the spirit is a question of the origin of the God. The question of the origin of the God is a question of the origin of the universe.

```

$IBSYS
$JOB          735003  WAVELENGTHS-PARTITION FUNCTIONS XY2 NON-LINEAR
$IBJOB H2TE     NODECK
$IBFTC PARRAT  NOLIST
      DIMENSION T(20), W(3), XM(10), UX(10), A(3,10), AL(3,10),
1  WW(3,10), CWW(3,10), U(140), PR(20), ZM(10)
      T(1)=0.
      DO 28 I=2,11
28  T(I)=T(I-1)+10.
      DO 29 I=12,20
29  T(I)=T(I-1)+100.
      W(1)=2047.
      W(2)=860.79
      W(3)=2133.
      READ (5,1) N
      DO 30 I=1,N
30  READ (5,2) ZM(I)
      YM=1.008
      C1=2.5425591*10.**5
      C2=0.23362281*10.**5
      C3=0.13090571*10.**5
      C4=-0.12*10.**5
      DO 31 I=1,N
      XM(I)=ZM(I)/(6.025*10.**3)
      UX(I)=((6.025*10.**3)/YM)+(1./XM(I))
      A(1,I)=(UX(I)*((2.*C2)+C4+C1))-(4.*C3/XM(I))
      A(2,I)=(2.*C2*(C1+C4)-(4.*(C3**2)))*((UX(I)**2)-(1./XM(I)**2))
      A(3,I)=(C1-C4)*UX(I)
      AL(1,I)=(A(1,I)+((A(1,I)**2)-4.*A(2,I))*0.5)/2.
      AL(2,I)=(A(1,I)-((A(1,I)**2)-4.*A(2,I))*0.5)/2.
31  AL(3,I)=A(3,I)
      DO 36 K=1,3
      DO 36 L=1,N
      DO 35 I=1,3
      DO 35 J=1,N
35  WW(I,J)=(AL(I,J)/(4.*((22./7.))**2)*9.))**0.5
36  CWW(K,L)=(W(K)/WW(K,1))*WW(K,L)
      WRITE (6,19)
      WRITE (6,16) (WW(1,J),J=1,N)
      WRITE (6,20)
      WRITE (6,16) (WW(2,J),J=1,N)
      WRITE (6,21)
      WRITE (6,16) (WW(3,J),J=1,N)
      WRITE (6,23)
      WRITE (6,16) (CWW(1,J),J=1,N)
      WRITE (6,24)
      WRITE (6,16) (CWW(2,J),J=1,N)
      WRITE (6,25)
      WRITE (6,16) (CWW(3,J),J=1,N)

```



```

V=1.438828
DO 37 L=1,20
UU=V/(T(L)+273.)
U(L)=UU*CW(1,7)
U(L+20)=UU*CW(1,1)
U(L+40)=UU*CW(2,7)
U(L+60)=UU*CW(2,1)
U(L+80)=UU*CW(3,7)
37 U(L+100)=UU*CW(3,1)
E=2.71828
DEG1=1
DEG2=1
DEG3=1
DO 38 I=1,20
H=-U(I)/2.
B=-U(I+20)/2.
C=-U(I+60)/2.
D=-U(I+40)/2.
F=-U(I+100)/2.
G=-U(I+80)/2.
P=-U(I+140)/2.
Q=-U(I+120)/2.
38 PR(I)=((U(I+20)*(E**B)*(1.-(E**(-U(I))))))
1/(U(I)*(E**H)*(1.-(E**(-U(I+20))))))*DEG1
2*((U(I+60)*(E**C)*(1.-(E**(-U(I+40))))))
3/(U(I+40)*(E**D)*(1.-(E**(-U(I+60))))))*DEG2
4*((U(I+100)*(E**F)*(1.-(E**(-U(I+80))))))
5/(U(I+80)*(E**G)*(1.-(E**(-U(I+100))))))*DEG3
WRITE (6,27)
WRITE (6,16) (PR(I),I=1,20)
1 FORMAT (I4)
2 FORMAT (2X,F10.5)
16 FORMAT (2X,F15.8)
19 FORMAT (22H THESE ARE THE OMEGA1S)
20 FORMAT (22H THESE ARE THE OMEGA2S)
21 FORMAT (22H THESE ARE THE OMEGA3S)
23 FORMAT (32H THESE ARE THE CORRECTED OMEGA1S)
24 FORMAT (32H THESE ARE THE CORRECTED OMEGA2S)
25 FORMAT (32H THESE ARE THE CORRECTED OMEGA3S)
27 FORMAT (31H THESE ARE THE PARTITION RATIOS)
CALL EXIT
END
$ENTRY          PARRAT
8
129.94853
127.94649
125.94420
124.94460
123.94278
122.94368
121.94193
119.94288

```


G. Computation of force constants, apex angle, isotopic vibrational frequencies, and partition function ratios for pyramidal XY_3 molecules

This program uses the Wilson F-G matrix method for finding the force constants. The apex angle is determined using Herzberg's formula (see page A30). Once these are known, isotopic vibrational frequencies and thus the partition function ratios can be computed. These are computed for the range of temperatures defined in parts A and F.

In the program, the following must be supplied:

YM = mass of the Y atom (awu)

W(I) = observed fundamental vibration frequencies
in cm^{-1} in order

DEGI = degree of degeneracy of the I^{th} vibration

Isotopic masses are read in as data cards, so for example, if the partition function ratio of the molecule containing the first isotope to the molecule containing the seventh isotope is desired, the values, CWW(1,7), CWW(1,1), CWW(2,7), CWW(2,1), CWW(3,7), CWW(3,1), etc. are used to define successive U's. If the ratio of the fourth isotope to the fifth is desired, one would use CWW(1,5), CWW(1,4), CWW(2,5), CWW(2,4), CWW(3,5), CWW(3,4), etc. in that order designating the U's from U(1) U(140).

The first part of the paper discusses the importance of the
theoretical framework in the study of the
relationship between the variables. The second part
presents the empirical results of the study. The third part
discusses the implications of the findings for the
theory and practice. The fourth part concludes the paper.

The first part of the paper discusses the importance of the
theoretical framework in the study of the
relationship between the variables. The second part
presents the empirical results of the study. The third part
discusses the implications of the findings for the
theory and practice. The fourth part concludes the paper.

The first part of the paper discusses the importance of the
theoretical framework in the study of the
relationship between the variables. The second part
presents the empirical results of the study. The third part
discusses the implications of the findings for the
theory and practice. The fourth part concludes the paper.

The first part of the paper discusses the importance of the
theoretical framework in the study of the
relationship between the variables. The second part
presents the empirical results of the study. The third part
discusses the implications of the findings for the
theory and practice. The fourth part concludes the paper.

The first part of the paper discusses the importance of the
theoretical framework in the study of the
relationship between the variables. The second part
presents the empirical results of the study. The third part
discusses the implications of the findings for the
theory and practice. The fourth part concludes the paper.

The first part of the paper discusses the importance of the
theoretical framework in the study of the
relationship between the variables. The second part
presents the empirical results of the study. The third part
discusses the implications of the findings for the
theory and practice. The fourth part concludes the paper.

```

$IRSYS
$JOB          735003 FORCE CONSTANTS - PARTITION FUNCTIONS, PYRAMIDAL-XY3
$TIME        005,300
$IBJOB TEO4   NODECK
$IBFTC FORCON NOLIST
  DIMENSION W(4), XM(10), UX(10), AB(4), BETA(10),
1ALPHA(10), ANGLE(10), G11A(10), G12A(10), G22A(10),
2G11E(10), G12E(10), G22E(10), AA(10), BA(10), AE(10),
3BE(10), AL(4,10), WW(4,10), T(20), U(160), PR(20),
4CWW(4,10), RR(20), SR(20), COA(20), COM(20), HRA(20), HRM(20)
  DO 28 I=2,11
    T(I)=0.
28 T(I)=T(I-1)+10.
    DO 29 I=12,20
29 T(I)=T(I-1)+100.
    YM=16.
    UY=(6.025*10.**3)/16.
    W(1)=758.
    W(2)=364.
    W(3)=703.
    W(4)=326.
    READ (5,1) N
    DO 30 I=1,N
30 READ (5,2) XM(I)
    DO 31 I=1,N
31 UX(I)=(6.025*10.**3)/XM(I)
    DO 32 I=1,4
32 AB(I)=4.*((22./7.)**2)*9.*(W(I)**2)
    CA=AB(1)+AB(2)
    DA=AB(1)-AB(2)
    CE=AB(3)+AB(4)
    DE=AB(3)-AB(4)
    DO 33 I=1,N
    R=1.
    BETA(I)=ARCOS(1./((4.*(W(3)**2)*(W(4)**2)/((W(1)**2)*(W(2)**2)))
1+((3.*YM)-XM(I))/((3.*YM)*XM(I))**0.5
    ALPHA(I)=2.*(ARSIN(((3.**0.5)/2.)*SIN(BETA(I))))
    ANGLE(I)=ALPHA(I)*57.2958
    G11A(I)=UY+((1.+2.*COS(ALPHA(I)))*UX(I))
    G12A(I)=-(2.*(1.+2.*COS(ALPHA(I)))*(1.-COS(ALPHA(I)))*UX(I))
1/(R*SIN(ALPHA(I)))
    G22A(I)=(2.*(1.+2.*COS(ALPHA(I)))*(UY+2.*UX(I)
1*(1.-COS(ALPHA(I))))/(R**2)*(1.+COS(ALPHA(I))))
    G11E(I)=UY+(UX(I)*(1.-COS(ALPHA(I))))
    G12E(I)=-(1.*(1.-COS(ALPHA(I))**2)*UX(I))/(R*SIN(ALPHA(I)))
    G22E(I)=((2.+COS(ALPHA(I)))*UY+((1.-COS(ALPHA(I))**2)*UX(I))/
1((R**2)*(1.+COS(ALPHA(I))))

```



```

EA=(4.*CA*(G12A(1)**2)/G11A(1))-(4.*CA*G22A(1))
HA=(4.*(G22A(1)**2))-(4.*G22A(1)*(G12A(1)**2))/G11A(1)
PA=(CA**2)-(DA**2)
EE=(4.*CE*(G12E(1)**2)/G11E(1))-(4.*CE*G22E(1))
HE=(4.*(G22E(1)**2))-(4.*G22E(1)*(G12E(1)**2))/G11E(1)
PE=(CE**2)-(DE**2)
F22A=(-EA-((EA**2)-4.*HA*PA)**0.5)/(2.*HA)
F22E=(-EE-((EE**2)-4.*HE*PE)**0.5)/(2.*HE)
F11A=(CA-F22A*G22A(1))/G11A(1)
F11E=(CE-F22E*G22E(1))/G11E(1)
FRR=(F11A-F11E)/3.
FR=(F11A+2.*F11E)/3.
FALPHA=(F22A+2.*F22E)/(3.*(R**2))
FALPAL=(F22A-F22E)/(3.*(R**2))
AA(I)=(F11A*G11A(I))+(F22A*G22A(I))
BA(I)=(F11A*F22A)*(G11A(I)*G22A(I)-G12A(I)**2)
AE(I)=(F11E*G11E(I))+(F22E*G22E(I))
33 BE(I)=(F11E*F22E)*(G11E(I)*G22E(I)-G12E(I)**2)
DO 34 J=1,N
AL(1,J)=(AA(J)+((AA(J)**2)-4.*BA(J))**0.5)/2.
AL(2,J)=(AA(J)-((AA(J)**2)-4.*BA(J))**0.5)/2.
AL(3,J)=(AE(J)+((AE(J)**2)-4.*BE(J))**0.5)/2.
34 AL(4,J)=(AE(J)-((AE(J)**2)-4.*BE(J))**0.5)/2.
DO 36 K=1,4
DO 36 L=1,N
DO 35 I=1,4
DO 35 J=1,N
35 WW(I,J)=(AL(I,J)/(4.*((22./7.）**2)*9.))**0.5
36 CWW(K,L)=(W(K)/WW(K,1))*WW(K,L)
V=1.438828
DO 37 L=1,20
UU=V/(T(L)+273.)
U(L)=UU*CWW(1,7)
U(L+20)=UU*CWW(1,1)
U(L+40)=UU*CWW(2,7)
U(L+60)=UU*CWW(2,1)
U(L+80)=UU*CWW(3,7)
U(L+100)=UU*CWW(3,1)
U(L+120)=UU*CWW(4,7)
37 U(L+140)=UU*CWW(4,1)
E=2.71828
DO 38 I=1,20
A=-U(I)/2.
B=-U(I+20)/2.
C=-U(I+60)/2.
D=-U(I+40)/2.
F=-U(I+100)/2.
G=-U(I+80)/2.
P=-U(I+140)/2.
Q=-U(I+120)/2.

```

... (faint, mostly illegible text) ...

```

DEG1=1
DEG2=1
DEG3=2
DEG4=2
PR(I)=(U(I+20)*(E**B)*(1.-(E**(-U(I))))))
1/(U(I)*(E**A)*(1.-(E**(-U(I+20))))))**DEG1
2*((U(I+60)*(E**C)*(1.-(E**(-U(I+40))))))
3/(U(I+40)*(E**D)*(1.-(E**(-U(I+60))))))**DEG2
4*((U(I+100)*(E**F)*(1.-(E**(-U(I+80))))))
5/(U(I+80)*(E**G)*(1.-(E**(-U(I+100))))))**DEG3
6*((U(I+140)*(E**P)*(1.-(E**(-U(I+120))))))
7/(U(I+120)*(E**Q)*(1.-(E**(-U(I+140))))))**DEG4
WRITE (6,3)
WRITE (6,16) (G11A(I),I=1,N)
WRITE (6,4)
WRITE (6,16) (G12A(I),I=1,N)
WRITE (6,5)
WRITE (6,16) (G22A(I),I=1,N)
WRITE (6,6)
WRITE (6,16) CA
WRITE (6,7)
WRITE (6,16) DA
WRITE (6,8)
WRITE (6,16) F22A
WRITE (6,9)
WRITE (6,16) F11A
WRITE (6,10)
WRITE (6,16) F22E
WRITE (6,11)
WRITE (6,16) F11F
WRITE (6,12)
WRITE (6,16) FRR
WRITE (6,13)
WRITE (6,16) FR
WRITE (6,14)
WRITE (6,16) FALPHA
WRITE (6,15)
WRITE (6,16) FALPAL
WRITE (6,17)
WRITE (6,16) (ALPHA(I),I=1,N)
WRITE (6,18)
WRITE (6,16) (ANGLE(I),I=1,N)
WRITE (6,19)
WRITE (6,16) (WW(1,J),J=1,N)
WRITE (6,20)
WRITE (6,16) (WW(2,J),J=1,N)
WRITE (6,21)
WRITE (6,16) (WW(3,J),J=1,N)
WRITE (6,22)
WRITE (6,16) (WW(4,J),J=1,N)

```



```

WRITE (6,23)
WRITE (6,16) (CWW(1,J),J=1,N)
WRITE (6,24)
WRITE (6,16) (CWW(2,J),J=1,N)
WRITE (6,25)
WRITE (6,16) (CWW(3,J),J=1,N)
WRITE (6,26)
WRITE (6,16) (CWW(4,J),J=1,N)
WRITE (6,27)
WRITE (6,16) (PR(I),I=1,20)
1 FORMAT (I4)
2 FORMAT (2X,F10.5)
3 FORMAT (13H THIS IS G11A)
4 FORMAT (13H THIS IS G12A)
5 FORMAT (13H THIS IS G22A)
6 FORMAT (11H THIS IS CA)
7 FORMAT (11H THIS IS DA)
8 FORMAT (13H THIS IS F22A)
9 FORMAT (13H THIS IS F11A)
10 FORMAT (13H THIS IS F22E)
11 FORMAT (13H THIS IS F11E)
12 FORMAT (12H THIS IS FRR)
13 FORMAT (11H THIS IS FR)
14 FORMAT (15H THIS IS FALPHA)
15 FORMAT (16H THIS IS FALPALP)
16 FORMAT (2X,E14.8)
17 FORMAT (14H THIS IS ALPHA)
18 FORMAT (25H THIS IS ALPHA IN DEGREES)
19 FORMAT (22H THESE ARE THE OMEGA1S)
20 FORMAT (22H THESE ARE THE OMEGA2S)
21 FORMAT (22H THESE ARE THE OMEGA3S)
22 FORMAT (22H THESE ARE THE OMEGA4S)
23 FORMAT (32H 3HFSF ARE THE CORRECTED OMEGA1S)
24 FORMAT (32H THESE ARE THE CORRECTED OMEGA2S)
25 FORMAT (32H THESE ARE THE CORRECTED OMEGA3S)
26 FORMAT (32H THESE ARE THE CORRECTED OMEGA4S)
27 FORMAT (31H THESE ARE THE PARTITION RATIOS)
CALL EXIT
END

```

\$ENTRY	FORCON
---------	--------

8	
129.94853	
127.94649	
125.94420	
124.94460	
123.94278	
122.94368	
121.94193	
119.94288	

H. Computation of partition function ratios for general molecules

This program computes the partition function ratio for any molecule for which isotopic vibrational data is known, using equation 16, page 49. The isotopic vibrational frequencies (cm^{-1}) are read in as data cards. If Q_2/Q_1 is wanted where subscript 1 refers to the light molecule, and 2 refers to the heavy, the data cards are ordered as follows: ω_1 (light), ω_1 (heavy), ω_2 (light), ω_2 (heavy), etc.

The degeneracy of the frequency is indicated by DEGI corresponding to $\omega(I)$. For example, the degeneracy of ω_2 for SCTe is 2, thus DEG2 = 2 etc. Should there be more or less than three frequencies per isotopic molecule in the partition function ratio, the statements of PR(I) must be altered accordingly. The program for TeF_6 (page A65) is an example where two of the normal frequencies are altered by isotopic substitution; whereas, the program for TeO_3^- is an example where four of the normal frequencies are altered by isotopic substitution.


```

$IBSYS
$JOB          735003      PARTITION FUNCTION RATIOS
$IBJOB SCTE    NODECK
$IBFTC PARRAT  NOLIST
      DIMENSION T(20), W(6), PR(20), U(120)
      A=1.438828
      T(1)=0.
      DO 24 I=2,11
24    T(I)=T(I-1)+10.
      DO 25 I=12,20
25    T(I)=T(I-1)+100.
      READ (5,1) N
      DO 26 I=1,N
26    READ (5,2) W(I)
      K=0
      DO 28 J=1,N
      DO 27 I=1,20
      K=K+1
27    U(K)=(A/(T(I)+273.))*W(J)
28    CONTINUE
      E=2.71828
      DEG1=1
      DEG2=2
      DEG3=1
      DO 29 I=1,20
      H=-U(I)/2.
      B=-U(I+20)/2.
      C=-U(I+60)/2.
      D=-U(I+40)/2.
      F=-U(I+100)/2.
      G=-U(I+80)/2.
      P=-U(I+140)/2.
      Q=-U(I+120)/2.
29    PR(I)=((U(I+20)*(E**B)*(1.-(E**(-U(I))))))
      1/(U(I)*(E**H)*(1.-(E**(-U(I+20))))))*DEG1
      2*((U(I+60)*(E**C)*(1.-(E**(-U(I+40))))))
      3/(U(I+40)*(E**D)*(1.-(E**(-U(I+60))))))*DEG2
      4*((U(I+100)*(E**F)*(1.-(E**(-U(I+80))))))
      5/(U(I+80)*(E**G)*(1.-(E**(-U(I+100))))))*DEG3
      DO 36 I=1,20
36    WRITE (6,8) PR(I)
      1 FORMAT(I4)
      2 FORMAT(2X,F7.2)
      8 FORMAT (2X,F15.8)
      CALL EXIT
      END
$ENTRY          PARRAT
6
1347.4004
1347.0
340.0
337.0
426.3703
423.0

```



```

$IBSYS
$JOB          735003      PARTITION FUNCTION RATIOS
$IBJOB TEF6      NODECK
$IBFTC PARRAT  NOLIST
      DIMENSION T(20), W(6), PR(20), U(120)
      A=1.438828
      T(1)=0.
      DO 24 I=2,11
24    T(I)=T(I-1)+10.
      DO 25 I=12,20
25    T(I)=T(I-1)+100.
      READ (5,1) N
      DO 26 I=1,N
26    READ (5,2) W(I)
      K=0
      DO 28 J=1,N
      DO 27 I=1,20
      K=K+1
27    U(K)=(A/(T(I)+273.))*W(J)
28    CONTINUE
      E=2.71828
      DEG1=3
      DEG2=3
      DO 29 I=1,20
      H=-U(I)/2.
      B=-U(I+20)/2.
      C=-U(I+60)/2.
      D=-U(I+40)/2.
      F=-U(I+100)/2.
      G=-U(I+80)/2.
29    PR(I)=(((U(I+20)*(F**B)*(1.-(E**(-U(I))))))
      1/(U(I)*(E**H)*(1.-(E**(-U(I+20))))))*DEG1
      2*((U(I+60)*(E**C)*(1.-(E**(-U(I+40))))))
      3/(U(I+40)*(E**D)*(1.-(E**(-U(I+60))))))*DEG2
      DO 36 I=1,20
36    WRITE (6,8) PR(I)
      1 FORMAT(I4)
      2 FORMAT(2X,F7.2)
      8 FORMAT (2X,E15.8)
      CALL EXIT
      END
$ENTRY          PARRAT
4
756.92
751.0
329.38
327.0

```




B29831